

## FIGURE INDEX

2.1	Relaxation mechanisms contributing to the static relative dielectric permittivity of liquid media along with corresponding indicative characteristic timescales. . . . .	11
2.2	Hypothetical processes defining the real solvation free energy of an ion. . . . .	19
2.3	Experimentally-elusive quantities required for the evaluation of intrinsic single-ion solvation free energies. . . . .	32
2.4	Fundamental experimental problems in the determination of thermodynamic parameters for processes involving ions. . . . .	34
3.1	Four basic choices defining a molecular model. . . . .	41
3.2	Three main types of theoretical models applicable to the evaluation of single-ion solvation free energies, as well as derivative thermodynamic solvation parameters. . . . .	43
3.3	Types of boundary conditions and electrostatic schemes commonly employed in classical atomistic simulations. . . . .	69
3.4	Common types of non-polarizable water models used in atomistic simulations. . . . .	81
3.5	Intrinsic electric potential at the center of an uncharged cavity within a pure liquid recalculated using P- or M-summation. . .	113
3.6	External or internal intrinsic bulk electric potentials calculated based on a spherical region using P- or M-averaging. . . . .	118
3.7	Air-liquid interfacial potential of a planar slab of a pure liquid calculated using P- or M-integration. . . . .	123
3.8	Solvent structuring and polarization around an ion as obtained from atomistic simulations. . . . .	132
4.1	Thermodynamic reactions relevant to ionic solvation. . . . .	197
4.2	Thermodynamic cycles relevant to ionic solvation. . . . .	200
4.3	Schematic illustrations regarding the properties of a system of phases at equilibrium. . . . .	216
4.4	Schematic classification of different types of phases in terms of their (di)electric properties. . . . .	225

4.5	Results of illustrative two-dimensional electrostatics calculations involving square or rectangular conductors surrounded by vacuum. . . . .	240
4.6	Results of illustrative two-dimensional electrostatics calculations involving square or rectangular conductors surrounded by vacuum. . . . .	242
4.7	Results of illustrative two-dimensional electrostatics calculations involving square conductors surrounded by vacuum. . . .	244
4.8	Definitions of the lateral and frontal Volta and surface potentials and potential differences for conducting phases. . . . .	249
4.9	Definition of the standard vacuum, real and intrinsic absolute potentials of an electrode. . . . .	265
4.10	Illustration of the connection between real and intrinsic absolute electrode potentials and corresponding real and intrinsic single-ion solvation free energies. . . . .	271
4.11	Potentials in a hypothetical fluid of hard spheres bearing isotropic quadrupole charge distributions. . . . .	278
4.12	External and internal Galvani potentials within a spherical sample of non-polarizable liquid argon. . . . .	280
4.13	Analogues of the SPC water model with altered charge distributions. . . . .	287
4.14	Schematic representation of an electrochemical equilibrium measurement. . . . .	291
4.15	Example of a Galvanic cell. . . . .	295
4.16	Schematic representation of Voltaic cell measurements. . . . .	301
4.17	Practical implementation of the Kenrick cell. . . . .	304
4.18	Summary of the relationships connecting key quantities related to single-ion solvation. . . . .	321
5.1	Available sets of effective ionic radii for the alkali and halide ions. . . . .	339
5.2	Standard electrode (redox) potentials and first isothermal temperature derivatives for the alkali and halide element-ion couples in water, along with contributions of more fundamental thermodynamic parameters to these quantities. . . . .	348
5.3	Standard thermodynamic parameters of dissolution for the alkali-halide salts in water. . . . .	362
5.4	Available estimates for the real absolute potential $\hat{V}_H^\ominus$ of the reference hydrogen electrode in water. . . . .	404
5.5	Available estimates for the real single-ion solvation free energy $\hat{G}_H^\ominus$ of the proton in water. . . . .	409
5.6	Available estimates for the standard air-liquid interfacial potential $\chi_{wat}^\ominus$ of water. . . . .	435

5.7	Estimates for the standard intrinsic hydration parameters of the proton and intrinsic partial molar variables of the aqueous proton. . . . .	461
5.8	Recommended data for the single-ion intrinsic hydration parameters of the gas-phase proton and alkali and halide ions, and the single-ion intrinsic partial molar variables of the aqueous proton and alkali and halide ions. . . . .	479
6.1	Correction terms to raw ionic solvation free energies calculated using atomistic simulations. . . . .	499
6.2	Illustration of the ion sizes corresponding to the three optimized ion-water Lennard-Jones interaction parameter sets. . . .	521
6.3	Properties that could in principle be used for the (in)validation of a specific ion-water (and ion-ion) Lennard-Jones interaction parameter set. . . . .	524
6.5	Thermodynamic cycles involved in the quantum-mechanical calculation of single-ion solvation free energies using quasi-chemical theory. . . . .	533