

LEVEL-SET PERCOLATION FOR THE GAUSSIAN FREE FIELD ON A TRANSIENT TREE

Angelo Abächerli and Alain-Sol Sznitman

Abstract

We investigate level-set percolation of the Gaussian free field on transient trees, for instance on super-critical Galton-Watson trees conditioned on non-extinction. Recently developed Dynkin-type isomorphism theorems provide a comparison with percolation of the vacant set of random interacements, which is more tractable in the case of trees. If h_* and u_* denote the respective (non-negative) critical values of level-set percolation of the Gaussian free field and of the vacant set of random interacements, we show here that $h_* < \sqrt{2u_*}$ in a broad enough set-up, but provide an example where $0 = h_* = u_*$ occurs. We also obtain some sufficient conditions ensuring that $h_* > 0$.

Résumé

Nous étudions la percolation de niveau pour le champ libre gaussien sur des arbres transients, par exemple sur des arbres de Galton-Watson surcritiques conditionnés à survivre. Des théorèmes de type isomorphisme de Dynkin récemment obtenus offrent un outil de comparaison avec la percolation de l'ensemble vacant pour les entrelacs aléatoires, qui se trouve être plus simple à étudier dans le cas des arbres. Si h_* et u_* désignent les valeurs critiques respectives de la percolation de niveau du champ libre gaussien, et de l'ensemble vacant des entrelacs aléatoires, nous montrons dans un cadre assez général que $h_* < \sqrt{2u_*}$, mais présentons un exemple pour lequel on a les égalités $0 = h_* = u_*$. Nous obtenons aussi des conditions suffisantes qui impliquent que $h_* > 0$.

Departement Mathematik
ETH Zürich
CH-8092 Zürich
Switzerland

0 Introduction

In this work we investigate level-set percolation of the Gaussian free field on a transient tree. Recently, over the last couple of years, various versions of Dynkin-type isomorphism theorems have related Gaussian free fields to random interacements, see for instance [9], [14], [15], [16], [17], [21]. They have fostered an interplay between level-set percolation of the Gaussian free field and percolation of the vacant set of random interacements. In the case of transient trees, the vacant cluster of random interacements at a given site can be expressed in terms of Bernoulli site percolation, see [20]. This makes percolation of the vacant set of random interacements more tractable, in particular with the help of the methods developed in [10], [11], [12]. In view of the above mentioned interplay, this feature raises the hope of gaining further insight into the more intricate level-set percolation of the Gaussian free field on a transient tree. This strategy was implemented in [17] in the case of $(d+1)$ -regular trees when $d \geq 2$. In particular, it was shown there that $0 < h_* < \sqrt{2u_*}$, if h_* and u_* stand for the respective critical values of level-set percolation of the Gaussian free field and of percolation of the vacant set of random interacements. Here, we resume this approach in the broader context of transient trees, in particular for super-critical Galton-Watson trees conditioned on non-extinction. Whereas we provide an example showing that $0 = h_* = \sqrt{2u_*}$ may occur, we prove under rather general assumptions that $h_* < \sqrt{2u_*}$, and derive sufficient conditions ensuring that $h_* > 0$.

Let us now describe the set-up and our results in more detail. We consider a locally finite tree (that is, a locally finite connected graph without loops) with vertex set T , such that each edge has unit weight and the corresponding weighted graph is transient. The discrete time random walk on T , when located in x , jumps to any given neighbor with probability $\deg(x)^{-1}$, where $\deg(x)$ stands for the degree of x . We write P_x for the canonical law of the walk starting in x , E_x for the corresponding expectation and $(X_k)_{k \geq 0}$ for the walk. The Green function is symmetric, positive, and equals

$$(0.1) \quad g(x, y) = \frac{1}{\deg(y)} E_x \left[\sum_{k=0}^{\infty} 1\{X_k = y\} \right], \text{ for } x, y \in T.$$

We write \mathbb{P}^G for the canonical law on \mathbb{R}^T of the Gaussian free field on T , and denote by $(\varphi_x)_{x \in T}$ the canonical field, so that under \mathbb{P}^G

$$(0.2) \quad (\varphi_x)_{x \in T} \text{ is a centered Gaussian field with covariance } g(\cdot, \cdot).$$

The critical value of the level-set percolation of φ is defined as

$$(0.3) \quad h_* = \inf\{h \in \mathbb{R}; \mathbb{P}^G\text{-a.s., all connected components of } \{\varphi \geq h\} \text{ are finite}\}$$

(here $\{\varphi \geq h\} = \{x \in T; \varphi_x \geq h\}$ and $\inf \phi = \infty$).

By a general argument of [2], recalled in the Appendix, one knows that

$$(0.4) \quad 0 \leq h_* \leq \infty.$$

Further, given $u \geq 0$, we consider the vacant set \mathcal{V}^u of random interacements at level u . This random subset of T is governed by a probability \mathbb{P}^I (see (1.32)) and \mathcal{V}^u becomes thinner as u increases. The critical value for the percolation of \mathcal{V}^u is defined as

$$(0.5) \quad u_* = \inf\{u \geq 0; \mathbb{P}^I\text{-a.s., all connected components of } \mathcal{V}^u \text{ are finite}\}.$$

To describe our results we introduce some base point x_0 of T , and define for any x in T , the sub-tree T_x of descendants of x , consisting of those y in T for which the geodesic path between x_0 and y goes through x (see the beginning of Section 1). We then write, see (1.4),

$$(0.6) \quad R_x^\infty = \text{the effective resistance between } x \text{ and } \infty \text{ in } T_x.$$

As an application of the cable graph methods initiated in [9], we show in Corollary 2.3 of Section 2 that when

$$(0.7) \quad \text{for some } A > 0, \text{ the (deterministic) set } \{x \in T; R_x^\infty > A\} \\ \text{only has finite components,}$$

then one has

$$(0.8) \quad 0 \leq h_* \leq \sqrt{2u_*}.$$

We also present in Remark 2.4 2) an example where

$$(0.9) \quad 0 = h_* = \sqrt{2u_*}.$$

As an aside one may wonder whether $\mathbb{P}^G[x_0 \xrightarrow{\varphi \geq 0} \infty] > 0$ holds under (0.7). This issue is linked to the geometry of the sign-clusters of the Gaussian free field on the cable system, see Remark 2.4 3).

Getting hold of strict inequalities strengthening (0.8) is more delicate. We provide in Theorem 3.4 a rather general sufficient condition, which ensures that $h_* < \sqrt{2u_*}$. This result comes as an application of the special coupling between random interacements and the Gaussian free field, which appears in Corollary 2.3 and was constructed in [17] as a refinement of [9]. More precisely, we show in Theorem 3.4 that when $0 < u_* < \infty$, and conditions (3.1) and (3.2) hold, that is, for some $A, B, M, \delta > 0$, for all distant vertices x in T having an infinite line of descent

$$(0.10) \quad \sum_{y \in (x_0, x)} 1\{R_y^\infty \leq A, d_{y^-} \leq M\} \geq \delta|x|,$$

$$(0.11) \quad \sum_{y \in (x_0, x]} \frac{1}{R_y^\infty(1 + R_y^\infty)} \leq B|x|$$

(with $|x|$ the distance of x to x_0 , and y^- the parent of y , see the beginning of Section 1 for notation), then, one has

$$(0.12) \quad h_* < \sqrt{2u_*}.$$

Importantly, to take advantage of the above mentioned special coupling, we show in Proposition 3.2 that (0.10) implies an exponential decay in $|x|$ of $\mathbb{P}^G[x_0 \xrightarrow{\varphi \geq 0} x]$. The proof is based on an idea of “entropic repulsion” in the spirit of [7], p. 13, 14.

In Proposition 4.2 we give a sufficient condition for $h_* > 0$, namely the existence of an infinite binary sub-tree of sites having uniformly bounded degree. As an application of Theorem 3.4 and Proposition 4.2, we see for instance that

$$(0.13) \quad 0 < h_* < \sqrt{2u_*} < \infty, \text{ when } T \text{ has bounded degree and outside a} \\ \text{finite subset of } T, \text{ each site has degree at least 3.}$$

Incidentally, in the case of \mathbb{Z}^d , $d \geq 3$, the inequality $h_* \leq \sqrt{2u_*}$ is known, see [9], but the strict inequality $h_* < \sqrt{2u_*}$ is presently open, and $h_* > 0$ is only known when d is sufficiently large, see [5], [13].

Our results also apply to typical realizations of super-critical Galton-Watson trees conditioned on non-extinction. In this case, one knows from [19] that u_* is deterministic and $0 < u_* < \infty$. There is even a reasonably explicit formula characterizing u_* , which is recalled in (5.4). We show in Lemma 5.1 that h_* is deterministic as well. In the more challenging Proposition 5.2 we show that (3.1) (or (0.10)) holds almost surely on non-extinction. In particular, as by-product, we deduce that

$$(0.14) \quad \begin{aligned} &0 \leq h_* \leq \sqrt{2u_*} < \infty, \text{ and almost surely on non-extinction} \\ &\mathbb{P}^G[x_0 \xrightarrow{\varphi \geq 0} x] \text{ has exponential decay in } |x|. \end{aligned}$$

When the offspring distribution has in addition some finite exponential moment (used to check (3.2)) we show in Theorem 5.4 that

$$(0.15) \quad h_* < \sqrt{2u_*}.$$

We also provide a sufficient condition for $h_* > 0$ in Theorem 5.5. We show in Theorem 5.5 that $h_* > 0$ when the offspring distribution has mean $m > 2$. Whereas Proposition 4.2 relies on the existence of an infinite binary sub-tree of sites having uniformly bounded degree, Theorem 5.5 follows a strategy in the spirit of Tassy [19] for random interacements on Galton-Watson trees, but the situation is more complicated in the case of the Gaussian free field. One can naturally wonder whether $h_* > 0$ holds generally when $m > 1$, see also Remark 5.6 .

We now explain the organization of this article. In Section 1 we introduce further notation and recall various facts concerning the Gaussian free field and random interacements. In Section 2 we consider the Gaussian free field $\tilde{\varphi}$ on the cable system attached to T . We introduce condition (2.3) (see also (0.7)), which enables us to prove in Proposition 2.2 that $\{\tilde{\varphi} > 0\}$ only has bounded components, and hence to apply the results of [9] and [17]. We show (0.8) in Corollary 2.3. In Section 3 we introduce the conditions (3.1), (3.2) (see (0.10), (0.11)), and show in Theorem 3.4 that together with the assumption $0 < u_* < \infty$, they imply that $h_* < \sqrt{2u_*}$. An important step established in Proposition 3.2 shows the exponential decay of $\mathbb{P}^G[x_0 \xrightarrow{\varphi \geq 0} x]$ under (3.1) (i.e. (0.10)). Section 4 provides a sufficient condition for $h_* > 0$ in Proposition 4.2 and the proof of (0.13) follows from Corollary 4.5, as explained in Remark 4.6. In Section 5 applications to super-critical Galton-Watson trees conditioned on non-extinction are discussed. The fact that h_* is deterministic (i.e. almost surely constant conditioned on non-extinction) appears in Lemma 5.1. The key Proposition 5.2 establishes that (3.1) holds almost surely on non-extinction, and (0.14) comes as a by-product, see also Theorem 5.4. Then, Proposition 5.3 shows that under the finiteness of some exponential moment of the offspring distribution, condition (3.2) holds almost surely on non-extinction. From that (0.15) readily follows in Theorem 5.4. In Theorem 5.5 we give sufficient conditions for $h_* > 0$. Finally, the Appendix contains the proof of the inequality $h_* \geq 0$ in the general set-up of transient weighted graphs, along similar arguments as in [2].

1 Some preliminaries

In this section we introduce further notation and collect useful results concerning transient trees, random walks, level-set percolation of the Gaussian free field, and percolation of the vacant set of random interlacements.

We consider a locally finite tree T with root x_0 such that each edge has unit weight and the resulting network is transient. We write $x \sim y$ when x and y are neighbors in T , we let $d(\cdot, \cdot)$ stand for the geodesic distance on the tree and $|x| = d(x, x_0)$ stand for the height of x in T . Given $U \subseteq T$, we let $\partial U = \{y \in T \setminus U; d(y, x) = 1, \text{ for some } x \in U\}$ and $\partial_i U = \{x \in U; d(y, x) = 1, \text{ for some } y \in T \setminus U\}$ respectively denote the outer and inner boundary of U . We let $|U|$ stand for the cardinality of U . For x in T we write $d(x, U) = \inf\{d(x, y); y \in U\}$ for the distance of x to U . Given x, y in T , we let $[x, y]$ stand for the collection of sites on the geodesic path from x to y . We also use the notation (x, y) , $[x, y)$, or $(x, y]$ when we exclude one or both endpoints. When $x \neq x_0$ we let x^- stand for the last point before x on the geodesic path from x_0 to x . Given $x \in T$, we denote by $d_x = |\{y \in T; y^- = x\}|$ the number of descendants of x , so that $\deg(x)$, the degree at x , equals d_{x_0} when $x = x_0$ and $d_x + 1$ when $x \neq x_0$. We let T_x stand for the sub-tree of descendants of x , i.e. consisting of those y in T for which x belongs to $[x_0, y]$. As far as dependence on the choice of the root is concerned, note that if a new root x'_0 is chosen, then T_x remains unchanged as soon as $x \notin [x_0, x'_0]$. Finally, a cut-set C separating x_0 from infinity (we will write cut-set for short) is a finite subset of $T \setminus \{x_0\}$, such that $x \neq y$ in C implies that $x \notin T_y$ (and $y \notin T_x$), and the connected component U_C of x_0 after deletion of the edges $\{x^-, x\}$, $x \in C$, is finite. We write $B_C = U_C \cup C$.

We now introduce some notation concerning simple random walk and potential theory on T . Given $U \subseteq T$, we write $T_U = \inf\{k \geq 0; X_k \notin U\}$ for the exit time of U , $H_U = \inf\{k \geq 0; X_k \in U\}$ for the entrance time in U , and $\tilde{H}_U = \inf\{k \geq 1; X_k \in U\}$ for the hitting time of U of the canonical walk $(X_k)_{k \geq 0}$ on T .

With similar notation as in (0.1), the Green function killed outside U is

$$(1.1) \quad g_U(x, y) = \frac{1}{\deg(y)} E_x \left[\sum_{0 \leq k < T_U} 1\{X_k = y\} \right], \text{ for } x, y \in T.$$

It is symmetric and vanishes when x or y does not belong to U . When $U = T$, we recover the Green function $g(x, y)$ from (0.1).

For K finite subset of T , the equilibrium measure of K is defined as

$$(1.2) \quad e_K(x) = \deg(x) P_x[\tilde{H}_K = \infty] 1_K(x), \text{ for } x \in T.$$

It is concentrated on the inner boundary of K and satisfies the identity

$$(1.3) \quad P_x[H_K < \infty] = \sum_y g(x, y) e_K(y), \text{ for } x \in T.$$

The total mass of e_K is the capacity $\text{cap}(K)$ of K .

As mentioned in the Introduction, an important quantity for x in T is the positive (possibly infinite) quantity

$$(1.4) \quad R_x^\infty = \text{the effective resistance between } x \text{ and } \infty \text{ in } T_x$$

(in particular $R_x^\infty = \infty$ when $|T_x| < \infty$, and R_x^∞ is the non-decreasing limit in N of the effective resistance in T_x between x and $\{x' \in T_x; d(x, x') = N\}$, when $|T_x| = \infty$).

As an aside, note that by the observation made above (1.1) moving the root x_0 to a different location x'_0 will only change finitely many of the R_x^∞ , $x \in T$. We then define

$$(1.5) \quad \alpha_x = \frac{R_x^\infty}{1 + R_x^\infty} \in (0, 1], \text{ for } x \in T,$$

as well as for $0 < \alpha \leq 1$ the operator

$$(1.6) \quad Q^\alpha f(a) = E^Y[f(\alpha a + \sqrt{\alpha} Y)], \text{ for } a \in \mathbb{R},$$

where Y stands for a standard normal variable, E^Y for the corresponding expectation, and f for a bounded measurable function. Note that for $\alpha = 1$, the above Q^α coincides with the Brownian transition kernel at time 1.

We now turn to the Gaussian free field φ on T . For $U \subseteq T$ we denote by σ_U the σ -algebra

$$(1.7) \quad \sigma_U = \sigma(\varphi_x, x \in U).$$

From the Markov property of the Gaussian free field, one knows that for x, y in T with $y^- = x$,

$$(1.8) \quad (\varphi_{y'} - P_{y'}[H_x < \infty] \varphi_x)_{y' \in T_y} \text{ is a centered Gaussian field with covariance } g_{U=T_y}(\cdot, \cdot) \text{ independent of } \sigma_{T \setminus T_y}.$$

The next lemma relates the objects we have now introduced, and will be recurrently used in this work ((1.15) will be used in the proof of Proposition 2.2 in Section 2).

Lemma 1.1. *For x in T , one has*

$$(1.9) \quad g(x, x) \leq R_x^\infty, \text{ with equality when } x = x_0,$$

$$(1.10) \quad \begin{cases} \text{i) } g(x, x) \geq 1/\deg(x), \\ \text{ii) } R_x^\infty \geq 1/d_x. \end{cases}$$

For x, y in T with $y^- = x$,

$$(1.11) \quad P_y[H_x < \infty] = \alpha_y, \quad P_y[H_x = \infty] = (1 + R_y^\infty)^{-1},$$

$$(1.12) \quad g_{U=T_y}(y, y) = \alpha_y,$$

and for any bounded measurable function f on \mathbb{R} one has

$$(1.13) \quad \mathbb{E}^G[f(\varphi_y) \mid \sigma_{T \setminus T_y}] = Q^{\alpha_y} f(\varphi_x).$$

When C is a cut-set, one has the identities

$$(1.14) \quad e_C(x) = \frac{1}{R_x^\infty} \text{ when } x \in C, \text{ and}$$

$$(1.15) \quad 1 = \sum_{x \in C} g(x_0, x) \frac{1}{R_x^\infty}.$$

Proof. The claims (1.9) and (1.10) follow from the fact that $g(x, x)$ is the effective resistance between x and infinity in T , whereas R_x^∞ is the effective resistance between x and infinity in T_x . As for (1.11), set $T'_y = \{x\} \cup T_y$, then the effective conductance between x and infinity in the sub-tree T'_y coincides with the escape probability $P_y[H_x = \infty]$, see also [12], above Proposition 17.26, so that $P_y[H_x = \infty] = (1 + R_y^\infty)^{-1}$ and $P_y[H_x < \infty] = \alpha_y$. Concerning (1.12), note that $g_{U=T_y}(y, y)$ coincides with the effective resistance between y and $\{x\} \cup \{\infty\}$ in T'_y , so that $g_{U=T_y}(y, y) = (1 + \frac{1}{R_y^\infty})^{-1} = \alpha_y$, whence (1.12).

We now turn to (1.13). By (1.8) we know that $\varphi_y - P_y[H_x < \infty] \varphi_x$ is a Gaussian variable with variance $g_{U=T_y}(y, y)$. By (1.11), (1.12), and the formula (1.6) defining Q^α the claim (1.13) readily follows. Concerning (1.14), we recall the notation B_C for C a cut-set, see above (1.1). One has the equality

$$(1.16) \quad e_{B_C} = e_C,$$

so that

$$(1.17) \quad e_C(x) \stackrel{(1.2)}{=} \sum_{y^-=x} P_y[H_x = \infty] \stackrel{(1.11)}{=} \sum_{y^-=x} (1 + R_y^\infty)^{-1} = \frac{1}{R_x^\infty}, \text{ for } x \in C.$$

Moreover, (1.15) is now the direct application of (1.3) (with the choice $x = x_0$, $K = B_C$) together with (1.14). This concludes the proof of Lemma 1.1. \square

Remark 1.2. Incidentally, when y_n , $0 \leq n < N$, with $N < \infty$ or $N = \infty$, is a finite or semi-infinite geodesic path in T moving away from the root x_0 , and $R_{y_n}^\infty = \infty$ for each $1 \leq n < N$, it follows from (1.13) and from the observation made below (1.6) that $(\varphi_{y_n})_{0 \leq n < N}$ under \mathbb{P}^G is distributed as a Brownian motion with the initial law $N(0, g(y_0, y_0))$, sampled at the integer times $0 \leq n < N$. \square

We now continue with level-set percolation of the Gaussian free field. Given x in T , and h in \mathbb{R} , we denote by $\{x \overset{\varphi \geq h}{\longleftrightarrow} \infty\}$ the event that the connected component of $\{\varphi \geq h\}$ containing x is infinite. If $\mathbb{P}^G[x \overset{\varphi \geq h}{\longleftrightarrow} \infty] > 0$ and y is neighbor of x , it is straightforward with (1.8) (where x plays the role of x_0) to infer that $\mathbb{P}^G[y \overset{\varphi \geq h}{\longleftrightarrow} \infty] > 0$ (one can also use the FKG-Inequality, see the Appendix of [7]). In other words, if $\mathbb{P}^G[x \overset{\varphi \geq h}{\longleftrightarrow} \infty]$ vanishes for some x in T , it vanishes for all x in T , and so we can express the critical value h_* defined in (0.3) as

$$(1.18) \quad h_* = \inf \{h \in \mathbb{R}; \mathbb{P}^G[x_0 \overset{\varphi \geq h}{\longleftrightarrow} \infty] = 0\} \text{ (with } x_0 \text{ the root)}.$$

By an argument of [2] one knows (actually, in the general set-up of transient weighted graphs, see Proposition A.2 of the Appendix) that

$$(1.19) \quad 0 \leq h_* \leq \infty.$$

Incidentally, in the case of \mathbb{Z}^d , $d \geq 3$, one knows that $h_* < \infty$ for all $d \geq 3$, see [2], [13], but $h_* > 0$ has only been proved when d is large enough, see [13], [5].

To further characterize h_* , we will now construct for each $h \in \mathbb{R}$ and x in T a $[0, 1]$ -valued function $q_{x,h}(\cdot)$, which is a “good version” of the conditional expectation $\mathbb{P}^G[x \xleftrightarrow{T_x, \varphi \geq h} \infty \mid \varphi_x = \cdot]$, where $\{x \xleftrightarrow{T_x, \varphi \geq h} \infty\}$ refers to the event that the connected component of x in $T_x \cap \{\varphi \geq h\}$ is finite.

For the definition we will now give, it is convenient to broaden the set-up, so that T is a tree with root x_0 , which is possibly recurrent or even finite. If T is recurrent then we set $R_x^\infty = \infty$ and $\alpha_x = 1$, for all $x \in T$.

For each $n \geq 0$, we write

$$(1.20) \quad T_n = \{x \in T; |x| = n\}, \quad B_n = \{x \in T; |x| \leq n\}$$

(so T_n is possibly empty, when T is finite).

Then, for each $n \geq 0$, $h \in \mathbb{R}$, $x \in B_n$, we define the functions $q_{x,h}^n(\cdot)$ by recursion towards the root x_0 , starting from the boundary T_n , via

$$(1.21) \quad \begin{cases} q_{x,h}^n(a) = 1_{(-\infty, h)}(a), \text{ for } x \in T_n \\ q_{x,h}^n(a) = 1_{(-\infty, h)}(a) + 1_{[h, \infty)}(a) \prod_{y^- = x} Q^{\alpha_y}(q_{y,h}^n)(a), \text{ for } |x| < n, \end{cases}$$

and an empty product (when x has no descendent) is understood as equal to 1.

Note that when T is finite and $n \geq 1$ such that $T_n = \emptyset$, then

$$(1.22) \quad q_{x,h}^n(\cdot) = 1, \text{ for all } x \in T.$$

Lemma 1.3. (*T a possibly finite or recurrent tree with root x_0*)

The functions $q_{x,h}^n(a)$, for $n \geq |x|$, are non-increasing in a , $[0, 1]$ -valued, equal to 1 on $(-\infty, h)$, with only possible discontinuity at h . For a fixed $x \in T$, they increase with $n \geq |x|$, and converge to a function $q_{x,h}(a)$ with similar properties, and such that

$$(1.23) \quad q_{x,h} = 1_{(-\infty, h)} + 1_{[h, \infty)} \prod_{y^- = x} Q^{\alpha_y}(q_{y,h}), \text{ for all } x \in T$$

(and an empty product is understood as equal to 1).

When T is finite, then

$$(1.24) \quad q_{x,h} = 1, \text{ for all } h \in \mathbb{R}, x \in T.$$

When T is transient, then for any $h \in \mathbb{R}$, $x \in T$, one has

$$(1.25) \quad q_{x,h}^n(\varphi_x) \stackrel{\mathbb{P}^G\text{-a.s.}}{=} \mathbb{P}^G[x \xleftrightarrow{T_x, \varphi \geq h} T_n \mid \varphi_x], \text{ for } n \geq |x|,$$

$$(1.26) \quad q_{x,h}(\varphi_x) \stackrel{\mathbb{P}^G\text{-a.s.}}{=} \mathbb{P}^G[x \xleftrightarrow{T_x, \varphi \geq h} \infty \mid \varphi_x].$$

In addition, one has the dichotomy

$$(1.27) \quad \begin{cases} \text{i) } q_{x_0, h} = 1 \text{ for } h > h_*, \\ \text{ii) } q_{x_0, h} \text{ is not identically 1 for } h < h_*. \end{cases}$$

Proof. From the definition of Q^α in (1.6) and the recursion from the boundary (1.21), it is immediate that $q_{x,h}^n(\cdot)$ are non-increasing, $[0, 1]$ -valued functions, equal to 1 on $(-\infty, h)$, with only possible discontinuity at h . When $x \in T_n$, $q_{x,h}^n(\cdot) \leq q_{x,h}^{n+1}(\cdot)$ by (1.21) and this gets propagated inside B_n by the recursion (1.21), so that $q_{x,h}^n(\cdot) \leq q_{x,h}^{n+1}(\cdot)$ for $x \in B_n$ (when $T_n = \phi$, actually (1.22) holds). Setting $q_{x,h}(a) = \lim_n \uparrow q_{x,h}^n(a)$, we obtain (1.23) from (1.21) by monotone convergence. It also follows that $q_{x,h}(\cdot)$ is non-increasing $[0, 1]$ -valued, with value 1 on $(-\infty, h)$ and only possible discontinuity at h (due to (1.23) and (1.6)). The claim (1.24) for finite T is immediate from (1.22).

Let us now assume that T is transient and prove (1.25). We fix n and use induction on $n - |x|$. When $x \in T_n$, then (1.25) is immediate from the first line of (1.21). When $|x| < n$, then one has the \mathbb{P}^G -a.s. equality

$$(1.28) \quad \mathbb{P}^G \left[x \xleftrightarrow{T_x, \varphi \geq h} T_n \mid \varphi_x \right] = 1_{(-\infty, h)}(\varphi_x) + 1_{[h, \infty)}(\varphi_x) \mathbb{P}^G \left[\bigcap_{y^- = x} \{y \xleftrightarrow{T_y, \varphi \geq h} T_n\} \mid \varphi_x \right].$$

If one first conditions on φ_x and φ_y , for $y^- = x$ it follows from the Markov property (1.8) and induction that we have \mathbb{P}^G -a.s.,

$$\mathbb{P}^G \left[\bigcap_{y^- = x} \{y \xleftrightarrow{T_y, \varphi \geq h} T_n\} \mid \varphi_x \right] = \mathbb{E}^G \left[\prod_{y^- = x} q_{y,h}^n(\varphi_y) \mid \varphi_x \right] \stackrel{(1.8), (1.13)}{=} \prod_{y^- = x} Q^{\alpha_y}(q_{y,h}^n)(\varphi_x).$$

Inserting this identity in the last expression of (1.28), and using (1.21), we see that (1.25) holds for x as well. This completes the proof of (1.25) by induction. The claim (1.26) readily follows by monotone convergence.

Finally, let us prove (1.27). We know that for $h > h_*$, $\mathbb{P}^G[x_0 \xleftrightarrow{\varphi \geq h} \infty] = 1$, cf. (1.18). By (1.26) we see that $q_{x_0,h}(\cdot) = 1$ almost everywhere and hence everywhere due to the fact that $q_{x_0,h}(\cdot)$ is non-increasing. This proves (1.27) i). On the other hand, for $h < h_*$, $\mathbb{P}^G[x_0 \xleftrightarrow{\varphi \geq h} \infty] < 1$, and by (1.26), $q_{x_0,h}(\cdot)$ is not identically 1, whence (1.27) ii). This completes the proof of Lemma 1.3. \square

Remark 1.4. Note that the recursion (1.21) used in the construction of the functions $q_{x,h}^n$ only involves the coefficients α_y , for $y \in T_x$. If we write $q_{x,h}^T$ for $q_{x,h}$, with $x \in T$ and $h \in \mathbb{R}$, to underline the dependence in T , it is straightforward to infer from (1.21) and the above observation that for all $x \in T$ and $h \in \mathbb{R}$

$$(1.29) \quad q_{x,h}^T = q_{x,h}^{T_x}.$$

This identity combined with (1.27) will be useful when proving that h_* is almost surely constant for a super-critical Galton-Watson tree conditioned on non-extinction. \square

We now return to the case where T is a transient tree with root x_0 , and consider the sub-tree (with same root x_0) of vertices with an infinite line of descent

$$(1.30) \quad T^\infty = \{x \in T; |T_x| = \infty\}.$$

Then, the connected components of $T \setminus T^\infty$ consist of finite sub-trees, and T^∞ is a transient tree with Green's function equal to the restriction of $g(\cdot, \cdot)$ to T^∞ , see for instance Proposition 1.11 of [18]. Thus, the law of $(\varphi_x)_{x \in T^\infty}$ under \mathbb{P}^G equals the law of the Gaussian free

field on T^∞ . Note also that for any $h \in \mathbb{R}$ one has $\{x_0 \xrightarrow{\varphi \geq h} \infty\} = \{x_0 \xrightarrow{T^\infty, \varphi \geq h} \infty\}$ (where this last notation refers to the event that the connected component of x_0 in $T^\infty \cap \{\varphi \geq h\}$ is infinite), so that with hopefully obvious notation

$$(1.31) \quad h_*(T) = h_*(T^\infty),$$

i.e. the critical values for level-set percolation of the Gaussian free field on T and on T^∞ coincide.

We now briefly turn to the topic of random interlacements on T and recall some facts concerning the percolation of the vacant set of random interlacements. We refer to the monographs [3] and [4] for further material and references. The vacant set of random interlacements at level $u \geq 0$ on T is a random subset \mathcal{V}^u of T , governed by a probability \mathbb{P}^I , with law characterized by

$$(1.32) \quad \mathbb{P}^I[\mathcal{V}^u \supseteq K] = \exp\{-u \operatorname{cap}(K)\}, \text{ for any finite } K \subseteq T$$

(with $\operatorname{cap}(K)$ the capacity of K , see below (1.3)).

As u increases, \mathcal{V}^u becomes thinner, and to classify the percolative properties of \mathcal{V}^u , one defines u_* as in (0.5). Actually, one has regardless of the choice of the base point x_0 , see Corollary 3.2 of [20],

$$(1.33) \quad u_* = \inf\{u \geq 0; \mathbb{P}^I[x_0 \xrightarrow{\mathcal{V}^u} \infty] = 0\} \in [0, \infty].$$

One also knows by Theorem 5.1 of [20] (the bounded degree assumption stated there can be removed) that

$$(1.34) \quad \begin{aligned} &\text{the connected component } \mathcal{C}^{\mathcal{V}^u}(x_0) \text{ of } x_0 \text{ in } \mathcal{V}^u \text{ has the same law as the} \\ &\text{open cluster of } x_0 \text{ in an independent site Bernoulli percolation on } T, \\ &\text{for which each site } x \in T \text{ is open with probability } p_{x,u}, \end{aligned}$$

where

$$(1.35) \quad \begin{cases} p_{x_0,u} = e^{-u \operatorname{cap}(\{x_0\})}, \text{ and for } x \neq x_0 \\ p_{x,u} = e^{-u \deg(x)} P_x[d(X_n, x_0) > d(x, x_0), \text{ for all } n > 0] \cdot P_x[d(X_n, x_0) \geq d(x, x_0), \text{ for all } n \geq 0] \end{cases}$$

Taking into account that $\operatorname{cap}(\{x_0\}) = g(x_0, x_0)^{-1}$ as well as (1.9), (1.11), we see that for $u \geq 0$,

$$(1.36) \quad p_{x_0,u} = e^{-\frac{u}{R_{x_0}^\infty}} \text{ and for } x \neq x_0, p_{x,u} = e^{-\frac{u}{R_x^\infty(1+R_x^\infty)}}.$$

Remark 1.5. If T^∞ stands for the sub-tree of vertices with an infinite line of descent, see (1.30), then with hopefully obvious notation

$$(1.37) \quad R_x^\infty(T) = R_x^\infty(T^\infty) \text{ for all } x \in T^\infty$$

(all components of $T \setminus T^\infty$ are finite, and (1.37) can be seen by replacing in the approximation of $R_x^\infty(T)$ below (1.4) the set $\{x' \in T_x; d(x, x') = N\}$ by the set $\{x' \in T_x^\infty;$

$d(x, x') = N\}$). In particular, in view of (1.36), we find that with similar notation as in (1.37) one has

$$(1.38) \quad p_{x,u}(T) = p_{x,u}(T^\infty) \text{ for all } x \in T^\infty,$$

and in view of (1.33), (1.34),

$$(1.39) \quad u_*(T) = u_*(T^\infty),$$

i.e. the critical values for the percolation of the vacant set of random interlacements on T and on T^∞ coincide. \square

Let us close this section by mentioning that percolation of \mathcal{V}^u can be re-expressed in terms of the transience of T endowed with certain weights. More precisely, if one introduces on the edges $e = \{x^-, x\}$, for $x \in T \setminus \{x_0\}$ the weights

$$(1.40) \quad c_u(e) = e^{-u \sum_{y \in (x_0, x]} \frac{1}{R_y^\infty(1+R_y^\infty)}} \left(1 - e^{-u \frac{1}{R_x^\infty(1+R_x^\infty)}}\right)^{-1},$$

then, when $u > 0$ and $R_x^\infty < \infty$ for each $x \in T$, one knows from Theorem 2.1 of [11], see also Corollary 5.25 of [12], and (1.34), (1.36) that

$$(1.41) \quad \mathbb{P}^I[x_0 \xrightarrow{\nu^u} \infty] > 0 \text{ if and only if } T \text{ endowed with the weights (1.40) is transient.}$$

2 Some consequences of the cable methodology

In this section we consider the Gaussian free field $\tilde{\varphi}$ on the cable system \tilde{T} attached to T . We use it to infer as an application of the results of [9] and [17] the inequality $h_* \leq \sqrt{2u_*}$, as well as a coupling, which relates the level sets of the Gaussian free field and the vacant set of random interlacements on T , see Corollary 2.3. This coupling will be the main tool in the next section to derive (under suitable assumptions) the strict inequality $h_* < \sqrt{2u_*}$. In Remark 2.4 2) we also provide an example where h_* and u_* vanish.

As in the previous section, T is a transient tree with base point x_0 , such that each edge has unit conductance. The cable tree \tilde{T} is obtained by attaching to each edge $e = \{x, y\}$ of the tree a compact interval with length $\frac{1}{2}$ and endpoints respectively identified to x and y . One defines on \tilde{T} a continuous diffusion behaving as a standard Brownian motion in the interior of each such segment. It has a continuous symmetric Green function $\tilde{g}(z, z')$, $z, z' \in \tilde{T}$ with respect to the Lebesgue measure on \tilde{T} , which extends the Green function $g(\cdot, \cdot)$ of the discrete time walk on T , see (0.1). We refer to Section 2 of [9], Section 2 of [6], and Section 3 of [21] for more details.

We now turn to the Gaussian free field on the cable tree \tilde{T} . On the canonical space $\tilde{\Omega}$ of continuous real-valued functions on \tilde{T} endowed with the σ -algebra generated by the canonical coordinates $\tilde{\varphi}_z$ (we also sometimes write $\tilde{\varphi}(z)$), $z \in \tilde{T}$, we denote by $\tilde{\mathbb{P}}^G$ the probability, with corresponding expectation $\tilde{\mathbb{E}}^G$, such that

$$(2.1) \quad \text{under } \tilde{\mathbb{P}}^G, (\tilde{\varphi}_z)_{z \in \tilde{T}} \text{ is a centered Gaussian field with covariance } \tilde{\mathbb{E}}^G[\tilde{\varphi}_z \tilde{\varphi}_{z'}] = \tilde{g}(z, z').$$

In particular, looking at the restriction of $\tilde{\varphi}$ to T , we see that

$$(2.2) \quad \text{the law of } (\tilde{\varphi}_x)_{x \in T} \text{ under } \tilde{\mathbb{P}}^G \text{ is equal to } \mathbb{P}^G.$$

An important issue in this context is to establish that $\tilde{\mathbb{P}}^G$ -a.s., $\{\tilde{\varphi} > 0\}$ only has bounded components in \tilde{T} . As shown in [9], see also (1.33) of [17], when this condition holds, then for $u > 0$ one can couple $\{\varphi > \sqrt{2u}\}$ and \mathcal{V}^u so that $\{\varphi > \sqrt{2u}\} \subseteq \mathcal{V}^u$, a.s.. It then follows that $h_* \leq \sqrt{2u_*}$.

We will introduce a condition, see (2.3), which implies the above condition, but also enables us to apply the results of Section 2 of [17] (see Corollary 2.5 and Remark 2.6 therein), and construct a strengthened coupling between $\{\varphi > \sqrt{2u}\}$ and \mathcal{V}^u , see (2.20) below.

We thus introduce the condition, see (1.4) for notation,

$$(2.3) \quad \text{for some } A > 0, \text{ the set } \{x \in T; R_x^\infty > A\} \text{ only has bounded components.}$$

Remark 2.1.

1) The condition (2.3) as we now explain is equivalent to the existence of a sequence of cut-sets C_n , $n \geq 1$, and $A > 0$, such that (see above (1.1) for notation)

$$(2.4) \quad \left\{ \begin{array}{l} \text{i) } B_{C_n} \subseteq U_{C_{n+1}} \text{ for each } n \geq 1 \\ \text{ii) } \sup_{n \geq 1} \sup_{x \in C_n} R_x^\infty \leq A. \end{array} \right.$$

Indeed, (2.4) readily implies (2.3). Conversely, when (2.3) holds, one defines U_1 consisting of x_0 and the points linked to x_0 by a path where $R_x^\infty > A$ prior to reaching x_0 , and sets $C_1 = \partial U_1$. By induction one then defines U_{n+1} as the union of U_n, C_n and the collection of points linked to C_n by a path where $R_x^\infty > A$ prior to reaching C_n , and sets $C_{n+1} = \partial U_{n+1}$. Then $B_{C_n} = U_n \cup C_n$, for each $n \geq 1$, and (2.4) holds.

Let us also mention that when C_n is a sequence of cut-sets as in (2.4), then by i)

$$(2.5) \quad B_{C_n} \subseteq B_{C_{n+1}} \text{ and } d(x_0, C_n) \geq n, \text{ for all } n \geq 1.$$

2) As a result of the observation above (1.5), condition (2.3) does not depend on the choice of the base point x_0 in T . \square

The main result established in this section comes in the next proposition. Its consequences appear in Corollary 2.3.

Proposition 2.2. *Assume that (2.3) holds, then*

$$(2.6) \quad \tilde{\mathbb{P}}^G\text{-a.s.}, \{\tilde{\varphi} > 0\} \text{ only has bounded components in } \tilde{T}.$$

Proof. For x, y in T , we write $[\widetilde{x}, \widetilde{y}]$ for the geodesic segment in \widetilde{T} between x and y . One has the following identities, which are consequences of the strong Markov property of $(\widetilde{\varphi}_z)_{z \in \widetilde{T}}$, see Lemma 3.1 and Proposition 5.2 of [9]: for $x \in T$,

$$(2.7) \quad \widetilde{\mathbb{P}}^G[\widetilde{\varphi} \text{ does not vanish on } [\widetilde{x}_0, \widetilde{x}]] = \frac{2}{\pi} \arcsin\left(\frac{g(x_0, x)}{\sqrt{g(x_0, x_0)g(x, x)}}\right),$$

$$(2.8) \quad \widetilde{E}^G[\widetilde{\varphi}_{x_0} \widetilde{\varphi}_x, \widetilde{\varphi} \text{ does not vanish on } [\widetilde{x}_0, \widetilde{x}]] = g(x_0, x)$$

(and the notation in (2.8) refers to the product with the indicator function of the event following the comma).

We consider $A > 0$ and a sequence of cut-sets C_n , $n \geq 1$, as in (2.4). We will first work under the additional assumption that

$$(2.9) \quad d_x = 1, \text{ for all } x \in C_n \text{ and } n \geq 1$$

(where d_x stands for the number of descendents of x in T , see the beginning of Section 1). We will then treat the general case. We thus assume (2.9) and define for each $n \geq 1$,

$$(2.10) \quad \widetilde{\mathcal{Z}}_n = \{x \in C_n; \widetilde{\varphi} > 0 \text{ on } [\widetilde{x}_0, \widetilde{x}]\}.$$

In what follows, constants will possibly depend on d_{x_0} , $R_{x_0}^\infty$, A , and will change from line to line. Additional dependence will appear in the notation. By (1.9), (1.10) and (2.4) ii) and (2.9), we see that

$$(2.11) \quad c \leq g(x, x) \leq c' \text{ for } x = x_0 \text{ and } x \in \bigcup_{n \geq 1} C_n.$$

Thus, by (2.7) and (2.8), we see that for $n \geq 1$,

$$(2.12) \quad \begin{aligned} \widetilde{\mathbb{E}}^G\left[\sum_{x \in C_n} (1 + \widetilde{\varphi}_{x_0} \widetilde{\varphi}_x) 1\{\widetilde{\varphi} > 0 \text{ on } [\widetilde{x}_0, \widetilde{x}]\}\right] &\leq c \sum_{x \in C_n} g(x_0, x) \\ &\stackrel{(2.4) \text{ ii)}}{\leq} c' \sum_{x \in C_n} g(x_0, x) \frac{1}{R_x^\infty} \stackrel{(1.15)}{=} c'. \end{aligned}$$

As a result, it follows from Fatou's lemma that

$$(2.13) \quad \widetilde{\mathbb{E}}^G\left[\liminf_n \sum_{x \in C_n} (1 + \widetilde{\varphi}_{x_0} \widetilde{\varphi}_x) 1\{\widetilde{\varphi} > 0 \text{ on } [\widetilde{x}_0, \widetilde{x}]\}\right] \leq c'.$$

This bound implies that the event

$$\bigcup_{L \geq 1} \limsup_n \left\{ |\widetilde{\mathcal{Z}}_n| \leq L, |\widetilde{\varphi}_{x_0}| \sum_{x \in \widetilde{\mathcal{Z}}_n} \widetilde{\varphi}_x \leq L \right\}$$

has full $\widetilde{\mathbb{P}}^G$ -probability. Since $|\widetilde{\varphi}_{x_0}| > 0$, $\widetilde{\mathbb{P}}^G$ -a.s., we find that

$$(2.14) \quad \widetilde{\mathbb{P}}^G\left[\bigcup_{M \geq 1} \limsup_n \left\{ |\widetilde{\mathcal{Z}}_n| \leq M \text{ and } \sum_{x \in \widetilde{\mathcal{Z}}_n} \varphi_x \leq M \right\}\right] = 1.$$

Note that for any $0 < \alpha \leq 1$, we have for any $0 \leq a \leq M$ (see (1.6) for notation)

$$(2.15) \quad \begin{aligned} Q^\alpha 1_{(-\infty, 0)}(a) &= P^Y[\alpha a + \sqrt{\alpha} Y < 0] = P^Y[Y < -\sqrt{\alpha} a] \\ &\stackrel{a \leq M, \alpha \leq 1}{\geq} P^Y[Y < -M] = c(M). \end{aligned}$$

We now introduce for $n, M \geq 1$ the event $A_{M,n} = \{\sum_{x \in \tilde{\mathcal{Z}}_n} (1 + \tilde{\varphi}_x) \leq M\}$ as well as the σ -algebra

$$(2.16) \quad \tilde{\mathcal{F}}_n = \sigma\left(\tilde{\varphi}_z, z \in \bigcup_{x \in C_n} [\widetilde{x_0}, x]\right).$$

It now follows from the Markov property of $\tilde{\varphi}$, see (1.8) of [17], that on $A_{M,n}$

$$(2.17) \quad \begin{aligned} \tilde{\mathbb{P}}^G[|\tilde{\mathcal{Z}}_{n+1}| = 0 \mid \tilde{\mathcal{F}}_n] &\geq \tilde{\mathbb{P}}^G[\tilde{\varphi}_y < 0, \text{ for all } y \in T \text{ with } y^- \in \tilde{\mathcal{Z}}_n \mid \tilde{\mathcal{F}}_n] \\ &= \prod_{x \in \tilde{\mathcal{Z}}_n} \prod_{y^- = x} Q^{\alpha_y}(1_{(-\infty, 0)})(\tilde{\varphi}_x) \stackrel{(2.9), (2.15)}{\geq} c(M)^M \text{ (on } A_{M,n}). \end{aligned}$$

By Borel-Cantelli's lemma it then follows that

$$\tilde{\mathbb{P}}^G\text{-a.s., on } \limsup_n A_{M,n}, |\tilde{\mathcal{Z}}_k| = 0 \text{ for large } k.$$

Since $\tilde{\mathbb{P}}^G[\bigcup_{M \geq 1} \limsup_n A_{M,n}] = 1$ by (2.14), we find that

$$\tilde{\mathbb{P}}^G\text{-a.s., } |\tilde{\mathcal{Z}}_n| = 0, \text{ for large } n.$$

We have thus shown that under (2.9)

$$(2.18) \quad \tilde{\mathbb{P}}^G\text{-a.s., the connected component of } x_0 \text{ in } \{\tilde{\varphi} > 0\} \text{ is bounded.}$$

We will now remove assumption (2.9). In essence, we use a scaling argument. We denote by T^* the tree with vertex set consisting of T and the mid-points of the intervals in \tilde{T} linking neighboring vertices in T , and edges of unit weight linking each mid-point to the two end-points of the interval where it lies. If \tilde{T}^* denotes the corresponding cable graph, there is a natural bijection s from \tilde{T}^* onto \tilde{T} , which, in essence, ‘‘scales by $\frac{1}{2}$ ’’ each interval linking neighbors in T^* . The effective resistance between two points in \tilde{T}^* is then twice the effective resistance between their images in \tilde{T} . Then, looking at Green functions, $\tilde{g}^*(z_1^*, z_2^*) = 2\tilde{g}(s(z_1^*), s(z_2^*))$, for z_1^*, z_2^* in \tilde{T}^* . It now follows that $(\sqrt{2}\tilde{\varphi}_{s(z^*)})_{z^* \in \tilde{T}^*}$ under $\tilde{\mathbb{P}}^G$ has the same law as the Gaussian free field on \tilde{T}^* .

Now consider a sequence $C_n \subseteq T$, $n \geq 1$, as in (2.4), and denote by $C_n^* \subseteq T^*$, the collection of mid-points x_* of the intervals attached to $\{x, x^-\}$, with $x \in C_n$, for $n \geq 1$. Note that $R_{x_*}^\infty(T^*) = 1 + 2R_x^\infty(T)$, for such x and x_* , and C_n^* , $n \geq 1$ is a sequence of cut-sets of T^* satisfying (2.4) with $1 + 2A$ in place of A . Moreover, (2.9) holds for C_n^* , $n \geq 1$. By (2.18) we see that $\tilde{\mathbb{P}}^G$ -a.s. the connected component of x_0 in $\{z^* \in \tilde{T}^*; \sqrt{2}\tilde{\varphi}_{s(z^*)} > 0\}$ is bounded. This proves that (2.18) holds under (2.3).

Now, as observed in Remark 2.1 2), (2.3) remains true for any choice of the base point x_0 . Hence, under (2.3), $\tilde{\mathbb{P}}^G$ -a.s. the connected components of all $x \in T$ in $\{\tilde{\varphi} > 0\}$ are bounded. This implies (2.6) and concludes the proof of Proposition 2.2. \square

We can now apply the results of [17].

Corollary 2.3. *Assume that (2.3) holds, then*

$$(2.19) \quad 0 \leq h_* \leq \sqrt{2u_*}.$$

Moreover, for any $u > 0$, one can couple independent copies $(\varphi_x)_{x \in T}$ and \mathcal{V}^u of the Gaussian free field on T and the vacant set of random interlacements at level u on T , with $(\eta_x)_{x \in T}$ a Gaussian free field on T , so that

$$(2.20) \quad \text{for all } B \subseteq (0, \infty), \{x \in T; \eta_x \in \sqrt{2u} + B\} \subseteq \{x \in T; \varphi_x \in B\} \cap \mathcal{V}^u.$$

Proof. Since $g(x, x) \leq R_x^\infty$, cf. (1.9), condition (2.3) ensures that (1.43) of [17] holds. Moreover, Proposition 2.2 shows that condition (1.32) of [17] holds as well. The claims follow from Corollary 2.5 and Remark 2.6 of [17]. Actually, (2.19) follows from (2.6) alone by the argument of [12], see also (1.33) of [17]. \square

Remark 2.4.

1) Note that an infinite self-avoiding path in T starting at the root x_0 only visits the sub-tree T^∞ of vertices with an infinite line of descent. Since $R_x^\infty(T) = R_x^\infty(T^\infty)$ for all $x \in T^\infty$, cf. (1.37), we thus see that

$$(2.21) \quad \text{condition (2.3) holds for } T \text{ if and only if (2.3) holds for } T^\infty.$$

2) As the present example shows, it is possible that all terms coincide in (2.19). For instance, consider a tree T with root x_0 , such that $d_x = 1 \vee |x|$, for all $x \in T$. This tree is transient (it contains a binary tree rooted at the descendents of x_0). As we now explain, for this tree one has

$$(2.22) \quad 0 = h_* = \sqrt{2u_*}.$$

To this end, note that for all $y \in T$, $R_y^\infty \leq R_{x_0}^\infty < \infty$. Thus, one finds that for all x in T (with c a positive constant changing from place to place)

$$\frac{1}{R_x^\infty} = \sum_{y^- = x} \frac{1}{1 + R_y^\infty} \geq d_x (1 + R_{x_0}^\infty)^{-1} \geq c|x|,$$

so that

$$(2.23) \quad \frac{1}{R_x^\infty(1 + R_x^\infty)} \geq c|x|.$$

If for $n \geq 1$ we set $C_n = \{x \in T; |x| = n\}$, we obtain a sequence of cut-sets of T with $|C_n| \leq n^n$, for which (2.4) holds. It then follows that $0 \leq h_* \leq \sqrt{2u_*}$. Moreover, for any $u > 0$ we have for large n

$$(2.24) \quad \sum_{x \in C_n} e^{-u \sum_{y \in (x_0, x]} \frac{1}{R_y^\infty(1 + R_y^\infty)}} \left(1 - e^{-\frac{u}{R_x^\infty(1 + R_x^\infty)}}\right)^{-1} \leq n^n e^{-uc'n^2} (1 - e^{-ucn})^{-1} \xrightarrow[n \rightarrow \infty]{} 0.$$

This implies that T endowed with the weights $c_u(e)$ in (1.40) is a recurrent weighted graph (see for instance Corollary 4.2 of [10]). By (1.41), this implies that $\mathbb{P}^I[x_0 \xrightarrow{v^u} \infty] = 0$, and therefore that $u_* \leq u$. Since $u > 0$ is arbitrary, we see that $u_* = 0$ and (2.22) follows.

3) In light of the above example one may still wonder whether $\mathbb{P}^G[x_0 \xrightarrow{\varphi \geq 0} \infty] > 0$ holds under (2.3). As we now briefly explain this issue is closely connected to a question concerning the geometry of the sign-clusters of $\tilde{\varphi}$ the Gaussian free field on the cable system.

We first explain how the positivity of $\mathbb{P}^G[x_0 \xrightarrow{\varphi \geq 0} \infty]$ can be expressed in terms of the sign clusters of $\tilde{\varphi}$. Observe that the law of $(\text{sign}(\varphi_x))_{x \in T}$ under \mathbb{P}^G can be generated by first considering the connected components of $\{|\tilde{\varphi}| > 0\}$ that meet T (they are $\tilde{\mathbb{P}}^G$ -a.s. bounded by (2.6)), and then by drawing independent symmetric random signs for each of these components. Such a representation can for instance be deduced from the strong Markov property of $\tilde{\varphi}$, combined with an exploration starting from x_0 of the successive components of $\{|\tilde{\varphi}| > 0\}$ that meet T , see also Lemma 3.1 of [9].

Then, denote by T' the random tree obtained by collapsing the sites of T that belong to a same component of $\{|\tilde{\varphi}| > 0\}$. From the above representation, the positivity of $\mathbb{P}^G[x_0 \xrightarrow{\varphi \geq 0} \infty]$ means that with positive $\tilde{\mathbb{P}}^G$ -measure, the Bernoulli site percolation with parameter $1/2$ on T' percolates (incidentally, this is indeed the case in the example in 2) above, due to the massive branching of T' in this example).

From the above and Theorem 6.2 of [10], see also Theorem 5.15 of [12], the positivity of $\mathbb{P}^G[x_0 \xrightarrow{\varphi \geq 0} \infty]$ now implies that with positive $\tilde{\mathbb{P}}^G$ -measure, the so-called branching number of T' (measuring the growth of T') is at least 2.

Thus, as a companion to the above question concerning the positivity of $\mathbb{P}^G[x_0 \xrightarrow{\varphi \geq 0} \infty]$ under (2.3), one may also wonder whether under (2.3) the branching number of the random tree T' is necessarily bigger or equal to 2.

□

3 A sufficient condition for $h_* < \sqrt{2u_*}$

In this section, we introduce two conditions, cf. (3.1), (3.2) below, and we show in Theorem 3.4 that when u_* is non-degenerate these conditions imply that $h_* < \sqrt{2u_*}$. An important step is contained in Proposition 3.2, where an exponential decay of the point to root connection probability in $\{\varphi \geq 0\}$ is derived. As in the previous section, T is a transient tree with root x_0 , and T^∞ stands for the sub-tree of vertices with an infinite line of descent, cf. (1.30).

We now introduce the two conditions mentioned above. The first condition states that

$$(3.1) \quad \text{there exists } A, M, \delta > 0 \text{ such that for large } n \text{ and all } x \in T^\infty \text{ with } |x| = n, \\ \sum_{y \in (x_0, x)} 1\{R_y^\infty \leq A, d_{y^-} \leq M\} \geq \delta n$$

(recall the beginning of Section 1 for notation). The second condition is:

(3.2) there exists $B > 0$ such that for large n and all $x \in T^\infty$ with $|x| = n$,

$$\sum_{y \in (x_0, x]} \frac{1}{R_y^\infty (1 + R_y^\infty)} \leq Bn.$$

Remark 3.1.

1) Note that when (3.1) holds, $\{y \in T, R_y^\infty > A\}$ cannot contain an infinite geodesic path in T , so that (3.1) implies (2.3). In particular Corollary 2.3 holds as a consequence of (3.1).

2) As a result of the observations made above (1.1) and below (1.4) concerning the effect of moving the location of the root, one sees that when (3.1) holds relative to x_0 , it will also hold relative to any different root x'_0 with possibly different $A', M', \delta' > 0$. The same observation applies to condition (3.2). \square

Our first main result consists in the derivation of an upper bound showing the exponential decay of $\mathbb{P}^G[x_0 \xrightarrow{\varphi \geq 0} x]$ for large x in T^∞ , when (3.1) holds. In essence we will use a strategy of “entropic repulsion” to prove this exponential decay, see [7], p. 13, 14, with however a special twist, see Remark 3.3.

Proposition 3.2. *Assume (3.1). Then, there exists $\kappa(A, M, \delta) > 0$ such that*

(3.3) *for large n and all $x \in T^\infty$ with $|x| = n$, $\mathbb{P}^G[x_0 \xrightarrow{\varphi \geq 0} x] \leq 2e^{-\kappa n}$.*

Proof. For $x \in T$ with $|x| = n$, we write x_0, x_1, \dots, x_n for the geodesic path in T from x_0 to x . Looking at (3.1), we see that when $x \in T^\infty$ and $|x|$ is large, one of the two sums corresponding to $y \in (x_0, x)$ with $y = x_k$, k even, or k odd, is at least $\frac{\delta}{2}|x|$. Hence, we can find $n_0 \geq 10$, such that for all $x \in T^\infty$ with $|x| \geq n_0$,

(3.4) there is a subset $I_x \subseteq [x_2, x)$ with $|I_x| \geq \frac{\delta}{3}n$, and $y \neq y' \in I_x \implies d(y, y') \geq 2$,
and for each $y \in I_x$, $R_y^\infty \leq A$ and $d_{y^-} \leq M$.

Further, note that for $x \in T^\infty$ with $|x| = n \geq n_0$

(3.5) for all $y \in I_x$ and $y' = (y^-)^-$ (so $|y'| = |y| - 2$), $R_{y'}^\infty \leq A + 2$.

Then, for x in T^∞ , with $|x| \geq n_0$, we define the subsets in $[x_0, x]$

(3.6) $J_x = I_x \cup \{(y^-)^-; y \in I_x\}$ and $K_x = \{y^-; y \in I_x\}$.

The sites in I_x are at mutual distance at least 2, and we thus see that for $x \in T^\infty$, with $|x| = n \geq n_0$

(3.7) $\left\{ \begin{array}{l} \text{i) } J_x \cap K_x = \phi, |J_x| \geq \frac{\delta}{3}n, |K_x| \geq \frac{\delta}{3}n, \\ \text{ii) } \text{for all } y \in J_x, R_y^\infty \leq A + 2, \\ \text{iii) } \text{for all } y \in K_x, \text{ both neighbors of } y \text{ in } [x_0, x] \text{ belong to } J_x, \\ \text{iv) } \text{for all } y \in K_x, d_y \leq M. \end{array} \right.$

We now use a strategy in the spirit of the proof of “entropic repulsion estimates” in p. 13, 14 of [7] to bound $\mathbb{P}^G[x_0 \xleftrightarrow{\varphi \geq 0} x]$.

We consider $x \in T^\infty$ with $|x| = n \geq n_0$ and introduce the event

$$(3.8) \quad F_x = \left\{ \text{for at least } \frac{\delta}{6} n \text{ sites } y \in K_x, \varphi_{y^-} \leq 1 \text{ and } \varphi_{y^+} \leq 1 \right\},$$

where y^+ stands for the only descendant of $y \in K_x$ in $[x_0, x]$ (so y^-, y^+ are the two neighbors of y in $[x_0, x]$). We will show that the probabilities of $\{x_0 \xleftrightarrow{\varphi \geq 0} x\} \cap F_x$ and of $\{x_0 \xleftrightarrow{\varphi \geq 0} x\} \setminus F_x$ decay exponentially in $|x|$, see (3.16) and (3.22). Note that by (3.7) iii) the event F_x only depends on the restrictions of φ to J_x :

$$(3.9) \quad F_x \in \sigma_{J_x} \text{ (see (1.7) for notation).}$$

By the Markov property of the Gaussian free field, we find that

$$(3.10) \quad \text{under } \mathbb{P}^G, \text{ conditionally on } \sigma_{J_x}, (\varphi_y)_{y \in K_x} \text{ are independent and distributed as } a_y + \xi_y, \text{ where}$$

$$(3.11) \quad a_y = E_y[H_{J_x} < \infty, \varphi(X_{H_{J_x}})], \text{ and}$$

$$(3.12) \quad \xi_y \text{ is a centered Gaussian variable with variance } g_{T \setminus \{y^-, y^+\}}(y, y) = \text{the effective resistance from } y \text{ to } \{y^-, y^+\} \cup \{\infty\} \\ \stackrel{(3.7) \text{ iv)}}{\geq} (1 + M)^{-1}.$$

We introduce the shorthand notation

$$(3.13) \quad A_x = \{\varphi_y \geq 0; \text{ for all } y \in [x_0, x]\} (= \{x_0 \xleftrightarrow{\varphi \geq 0} x\}).$$

Then, as an application of (3.10) - (3.12), we find that

$$(3.14) \quad \mathbb{P}^G[A_x] = \mathbb{P}^G[A_x \setminus F_x] + \mathbb{P}^G[A_x \cap F_x] \text{ with} \\ \mathbb{P}^G[A_x \cap F_x] \stackrel{(3.10)-(3.12)}{\leq} \mathbb{E}^G \left[F_x, \prod_{y \in K_x} P^{\xi_y}[a_y + \xi_y \geq 0] \right],$$

where ξ_y has the distribution from (3.12) under P^{ξ_y} .

Note that for each $y \in K_x$ for which $a_y \leq 1$ holds, we have

$$(3.15) \quad P^{\xi_y}[a_y + \xi_y \geq 0] \leq P^{\xi_y}[1 + \xi_y \geq 0] = 1 - P^{\xi_y}[\xi_y \leq -1] \\ \stackrel{(3.12)}{\leq} 1 - P^Y[Y \geq \sqrt{1 + M}], \text{ where } Y \text{ is } N(0, 1)\text{-distributed.}$$

By definition of the event F_x in (3.8), on F_x there are at least $\frac{\delta}{6} n$ sites $y \in K_x$ such that $a_y \leq 1$. Hence, we see that

$$(3.16) \quad \mathbb{P}^G[A_x \cap F_x] \leq (1 - P^Y[Y \geq \sqrt{1 + M}])^{\frac{\delta}{6} n}.$$

We will now bound $\mathbb{P}^G[A_x \setminus F_x]$, i.e. the first term in the right-hand side of the first line of (3.14). On $A_x \setminus F_x$, there are at least $|K_x| - \frac{\delta}{6}n \geq \frac{\delta}{6}n$ sites $y \in K_x$ where $\max\{\varphi_{y^-}, \varphi_{y^+}\} \geq 1$ (both y^-, y^+ belong to J_x , cf. (3.7) iii). Since on A_x , $\varphi \geq 0$ on $[x_0, x]$, we see that

$$(3.17) \quad A_x \setminus F_x \subseteq \left\{ \sum_{y \in J_x} \varphi_y \geq \frac{\delta}{12} n \right\}$$

(two ‘‘consecutive’’ y in K_x might share the same neighbor in J_y , whence the term $\frac{\delta}{12}n$).

Note that $\sum_{y \in J_x} \varphi_y$ is a centered Gaussian variable under \mathbb{P}^G , hence we have

$$(3.18) \quad \mathbb{P}^G[A_x \setminus F_x] \stackrel{(3.17)}{\leq} \mathbb{P}^G \left[\sum_{y \in J_x} \varphi_y \geq \frac{\delta}{12} n \right] \leq \exp \left\{ -\frac{1}{2} \left(\frac{\delta}{12} \right)^2 \frac{n^2}{\text{var} \left(\sum_{y \in J_x} \varphi_y \right)} \right\},$$

and we can express the variance in the last term as

$$(3.19) \quad \text{var} \left(\sum_{y \in J_x} \varphi_y \right) = \sum_{y \in J_x} g(y, y) + 2 \sum_{y < y' \text{ in } J_x} g(y, y')$$

$$\stackrel{(1.11)}{=} \sum_{y \in J_x} g(y, y) \left(1 + 2 \sum_{y' > y, y' \in J_x} \prod_{y < y'' \leq y'} \alpha_{y''} \right),$$

where the notation $y < y'$ means that $y \in [x_0, y')$ and $y < y'' \leq y'$ is defined in a similar fashion. Note that for $y'' \in J_x$, by (3.7) ii) we have $R_{y''}^\infty \leq A + 2$ and therefore

$$(3.20) \quad \alpha_{y''} \leq \alpha_A \stackrel{\text{def}}{=} \frac{A+2}{A+3}, \text{ for all } y'' \in J_x.$$

Since $\alpha_{y''} \leq 1$, we can restrict the product in the last term of (3.19) to the $y'' \in J_x$ and obtain the upper bound

$$(3.21) \quad \text{var} \left(\sum_{y \in J_x} \varphi_y \right) \leq \sum_{y \in J_x} g(y, y) \left(1 + 2 \sum_{\ell \geq 1} \alpha_A^\ell \right), \text{ and since } R_y^\infty \leq A + 2 \text{ on } J_x$$

$$\stackrel{(1.9)}{\leq} |J_x| (A + 2) \left(1 + 2 \frac{\alpha_A}{1 - \alpha_A} \right) \leq n(A + 2)(2A + 5).$$

Coming back to (3.18) we find

$$(3.22) \quad \mathbb{P}^G[A_x \setminus F_x] \leq \exp \left\{ -\frac{1}{2} \frac{\delta^2}{144} \frac{n}{(A+2)(2A+5)} \right\}.$$

Collecting (3.16) and (3.22), and coming back to (3.14), we see that for all $x \in T^\infty$ with $|x| = n \geq n_0$, we have

$$(3.23) \quad \mathbb{P}^G[x_0 \xleftrightarrow{\varphi \geq 0} x] \leq (1 - P^Y[Y \leq -\sqrt{1+M}])^{\frac{\delta}{6}n} + \exp \left\{ -\frac{1}{2} \frac{\delta^2}{144} \frac{n}{(A+2)(2A+5)} \right\}.$$

The claim (3.3) readily follows. \square

Remark 3.3. In the first inequality of (3.18), it is important that we bound $\mathbb{P}^G[A_x \setminus F_x]$ in terms of a deviation of $\sum_{J_x} \varphi_y$ and *not* of $\sum_{(x_0, x)} \varphi_y$. The point is the following. Whereas the variance of $\sum_{J_x} \varphi_y$ grows at most linearly in n , as crucially shown in (3.21), the variance of $\sum_{(x_0, x)} \varphi_y$ may grow faster than linearly in n , due to the presence of long stretches where, for instance, $d_y = 1$ and α_y is close to 1. A faster than linear growth of the variance would destroy the exponential decay we obtain in (3.22). \square

We now come to the main result of this section. The proof uses the coupling between the Gaussian free field and random interacements stated in (2.20) of Corollary 2.3.

Theorem 3.4. *Assume that $0 < u_* < \infty$ and (3.1), (3.2) hold, then*

$$(3.24) \quad (0 \leq) h_* < \sqrt{2u_*}.$$

Proof. In view of (1.31), (1.39) and (1.37), we can assume that $T = T^\infty$. With κ as in (3.3) and B as in (3.2), we consider

$$(3.25) \quad 0 < u < u_* \text{ where } u = u_* - \rho \text{ with } \rho = \min\left(\frac{u_*}{2}, \frac{\kappa}{8B}\right).$$

We will show that

$$(3.26) \quad h_* \leq \sqrt{2u} (< \sqrt{2u_*}),$$

and the claim (3.24) will follow.

We use the same notation $\mathcal{C}^{\mathcal{V}^u}(x_0)$ as in (1.34). Since $u < u_*$, the event

$$(3.27) \quad \mathcal{P}_{x_0, u} = \{x_0 \xrightarrow{\mathcal{V}^u} \infty\} = \{|\mathcal{C}^{\mathcal{V}^u}(x_0)| = \infty\} \text{ has positive } \mathbb{P}^I\text{-measure.}$$

On the event $\mathcal{P}_{x_0, u}$, $\mathcal{C}^{\mathcal{V}^u}(x_0)$ is an infinite sub-tree of T , rooted at x_0 . However, $\mathcal{C}^{\mathcal{V}^u}(x_0)$ is “thin”. Specifically, if we perform an independent Bernoulli site percolation on the tree $\mathcal{C}^{\mathcal{V}^u}(x_0)$ with parameter $p_{x, 2\rho}$, for $x \in \mathcal{C}^{\mathcal{V}^u}(x_0)$, in the notation of (1.35), (1.36), the resulting connected component of x_0 under the joint law of \mathcal{V}^u and the above Bernoulli percolation is that of the cluster $\mathcal{C}^{\mathcal{V}^{u+2\rho}}$. Since $u + 2\rho > u_*$, cf. (3.25), this cluster is a.s. finite. As a consequence,

$$(3.28) \quad \text{on an event } \tilde{\mathcal{P}}_{x_0, u} \subseteq \mathcal{P}_{x_0, u} \text{ with } \mathbb{P}^I(\mathcal{P}_{x_0, u} \setminus \tilde{\mathcal{P}}_{x_0, u}) = 0, \text{ a.s. for the above Bernoulli site percolation the open cluster of } x_0 \text{ in } \mathcal{C}^{\mathcal{V}^u}(x_0) \text{ is finite.}$$

Let us describe how the proof now proceeds. We will use the above statement expressing the thinness of $\mathcal{C}^{\mathcal{V}^u}(x_0)$ when $\tilde{\mathcal{P}}_{x_0, u}$ occurs, to construct a sequence of cut-sets C_n of $\mathcal{C}^{\mathcal{V}^u}(x_0)$ tending to infinity, along which $\sum_{x \in C_n} e^{-\frac{3}{4}\kappa|x|}$ tends to 0, see (3.32). By the exponential decay shown in (3.3) of Proposition 3.2 and a union bound, this will imply that on $\tilde{\mathcal{P}}_{x_0, u}$ the connected component of x_0 in $\{\varphi \geq 0\} \cap \mathcal{C}^{\mathcal{V}^u}(x_0)$ is \mathbb{P}^G -a.s. finite, see (3.33). With the help of the coupling in (2.20), it will then be a simple matter to infer that \mathbb{P}^G -a.s., the connected component of x_0 in $\{\varphi > \sqrt{2u}\}$ is finite and the claim (3.26) will follow. With this strategy in mind, we now proceed with the proof.

As mentioned in (1.41), by Corollary 5.25 of [12], we know that on $\tilde{\mathcal{P}}_{x_0, u}$,

$$(3.29) \quad \mathcal{C}^{\mathcal{V}^u}(x_0) \text{ endowed with the weights } c_{2\rho}(e), \text{ for } e = \{x^-, x\}, x \in \mathcal{C}^{\mathcal{V}^u}(x_0) \setminus \{x_0\},$$

is a recurrent network.

By Corollary 4.2 of [10], it then follows that on $\tilde{\mathcal{P}}_{x_0, u}$ for any summable sequence of positive numbers $w_m > 0$, $m \geq 1$, one can find a sequence of cut-sets $C_n, n \geq 1$, of $\mathcal{C}^{\mathcal{V}^u}(x_0)$, with $d(x_0, C_n) \xrightarrow[n]{n} \infty$, such that

$$(3.30) \quad \lim_n \sum_{x \in C_n} w_{|x|} e^{-2\rho \sum_{y \in (x_0, x]} \frac{1}{R_y^\infty(1+R_y^\infty)}} \left(1 - e^{-2\rho \frac{1}{R_x^\infty(1+R_x^\infty)}}\right)^{-1} = 0.$$

We apply this observation with the choice

$$(3.31) \quad w_m = e^{-\frac{\kappa}{2}m}, \quad m \geq 1 \text{ (and } \kappa \text{ as in (3.3))},$$

and consider the corresponding sequence of cut-sets $C_n, n \geq 1$ of $\mathcal{C}^{\mathcal{V}^u}(x_0)$. By (3.2) and (3.30) we see that on $\tilde{\mathcal{P}}_{x_0, u}$

$$(3.32) \quad \lim_n \sum_{x \in C_n} e^{-\frac{\kappa}{2}|x| - 2\rho B|x|} = 0 \text{ (record for later use that } \frac{\kappa}{2} + 2\rho B \stackrel{(3.25)}{\leq} \frac{3}{4}\kappa).$$

Thus, on $\tilde{\mathcal{P}}_{x_0, u}$, for large n ,

$$(3.33) \quad \mathbb{P}^G[x_0 \xrightarrow[\varphi \geq 0]{} \infty \text{ in } \mathcal{C}^{\mathcal{V}^u}(x_0)] \leq \sum_{x \in C_n} \mathbb{P}^G[x_0 \xrightarrow[\varphi \geq 0]{} x] \stackrel{(3.3)}{\leq} 2 \sum_{x \in C_n} e^{-\kappa|x|} \stackrel{(3.32)}{\xrightarrow[n \rightarrow \infty]{}} 0.$$

This shows that

$$(3.34) \quad \mathbb{E}^I[\mathcal{P}_{x_0, u}, \mathbb{P}^G[x_0 \xrightarrow[\varphi \geq 0]{} \infty \text{ in } \mathcal{C}^{\mathcal{V}^u}(x_0)]] = 0.$$

As observed in Remark 3.1 1), condition (3.1) implies that (2.20) holds. Choosing $B = (0, \infty)$ in (2.20), we see that

$$(3.35) \quad \begin{aligned} \mathbb{P}^G[x_0 \xrightarrow[\varphi > \sqrt{2u}]{} \infty] &\stackrel{(2.20)}{\leq} \mathbb{P}^I \otimes \mathbb{P}^G[x_0 \xrightarrow[\mathcal{V}^u \cap \{\varphi > 0\}]{} \infty] \\ &= E^I[\mathcal{P}_{x_0, u}, \mathbb{P}^G[x_0 \xrightarrow[\varphi \geq 0]{} \infty \text{ in } \mathcal{C}^{\mathcal{V}^u}(x_0)]] \stackrel{(3.34)}{=} 0. \end{aligned}$$

This proves that (3.26) holds and concludes the proof of the theorem. \square

We will later apply Theorem 3.4 in Section 5 when T is the typical realization of a super-critical Galton-Watson process conditioned on non-extinction, see Theorem 5.4.

4 A sufficient condition for $h_* > 0$

In this section we provide a sufficient condition which ensures that $h_* > 0$. The main result appears in Proposition 4.2. We provide a first application of the results of this and the previous section in Corollary 4.5 and in Remark 4.6.

We begin with some notation and a lemma that will be useful in the proof of Proposition 4.2. For $h \in \mathbb{R}$, we denote by π_h the multiplication operator by $1_{[h, \infty)}$, so that for f function on \mathbb{R} , one has

$$(4.1) \quad \pi_h f = 1_{[h, \infty)} f.$$

We recall the notation Q^α from (1.6).

Lemma 4.1. *If f is a bounded, non-decreasing, right-continuous function on \mathbb{R} , which vanishes on $(-\infty, 0)$, then*

$$(4.2) \quad \text{for any } 0 < \alpha \leq 1, \pi_0 Q^\alpha f \text{ has the same properties as noted above for } f,$$

$$(4.3) \quad \text{for any } a \in \mathbb{R}, \pi_0 Q^\alpha f(a) \text{ is a non-decreasing function of } \alpha \in (0, 1].$$

Proof. The claim (4.2) is immediate by direct inspection of (1.6). We now prove (4.3). We can find a positive measure with finite mass, supported on $[0, \infty)$, such that $f(a) = \mu([0, a])$, for $a \geq 0$ (and $f(a) = 0$, for $a < 0$). Hence, we see that for $a \in \mathbb{R}$,

$$(4.4) \quad \begin{aligned} Q^\alpha f(a) &\stackrel{(1.6)}{=} E^Y [f(a\alpha + \sqrt{\alpha} Y)] = E^Y \left[\int_{[0, \infty)} 1\{a\alpha + \sqrt{\alpha} Y \geq b\} d\mu(b) \right] \\ &= \int_{[0, \infty)} P^Y \left[Y \geq \frac{b}{\sqrt{\alpha}} - \sqrt{\alpha} a \right] d\mu(b). \end{aligned}$$

Now, when $a \geq 0$, $b \geq 0$, the function $\alpha > 0 \mapsto P^Y [Y \geq \frac{b}{\sqrt{\alpha}} - \sqrt{\alpha} a]$ is non-decreasing, and the claim (4.3) follows. \square

The main result of this section comes next.

Proposition 4.2. *Assume that the tree T contains an infinite binary sub-tree \bar{T} rooted at x_0 , such that for some $M > 0$,*

$$(4.5) \quad \sup_{x \in \bar{T}} d_x \leq M.$$

Then, T is transient and

$$(4.6) \quad h_* > 0.$$

Remark 4.3. The example in Remark 2.4 2), where one moves the root to its second neighbor, shows that the sole existence of an infinite binary tree rooted at x_0 does not guarantee that $h_* > 0$ (in this example we have $0 = h_* = \sqrt{2u_*}$). \square

Proof of Proposition 4.2: The transience of T is immediate. We introduce by analogy with (1.20), (1.21), $\overline{T}_n = \{x \in \overline{T}; |x| = n\} = T_n \cap \overline{T}$, $\overline{B}_n = \{x \in \overline{T}; |x| \leq n\} = B_n \cap \overline{T}$, as well as the function $\overline{q}_{x,h}^n$, for $n \geq 0$, $x \in \overline{B}_n$, $h \in \mathbb{R}$, via

$$(4.7) \quad \begin{cases} \overline{q}_{x,h}^n = 1_{(-\infty,h)}, & \text{for } x \in \overline{T}_n, \\ \overline{q}_{x,h}^n = 1_{(-\infty,h)} + 1_{[h,\infty)} \prod_{y^-=x,y \in \overline{T}} Q^{\overline{\alpha}}(\overline{q}_{y,h}^n), & \text{for } x \in \overline{B}_{n-1}, \end{cases}$$

where

$$(4.8) \quad \overline{\alpha} = \frac{1}{M+1} \geq \alpha_x = \frac{R_x^\infty}{1+R_x^\infty}, \text{ for } x \in \overline{T}$$

(since $R_x^\infty \geq \frac{1}{M}$, by (1.10) ii) and (4.5)).

As in Lemma 1.3, we see that for any $x \in \overline{T}$, $h \in \mathbb{R}$, the functions $\overline{q}_{x,h}^n$ increase in $n \geq |x|$ to a function $\overline{q}_{x,h}$, which is non-increasing, $[0, 1]$ -valued, equal to 1 on $(-\infty, h)$, with only possible discontinuity at h , and such that

$$(4.9) \quad \overline{q}_{x,h} = 1_{(-\infty,h)} + 1_{[h,\infty)} \prod_{y^-=x,y \in \overline{T}} Q^{\overline{\alpha}}(\overline{q}_{y,h})$$

(although we will not explicitly need it, it is also straightforward to see that $\overline{q}_{x,h} = \overline{q}_{x_0,h}$ for all $x \in \overline{T}$).

As we now explain, when $h \geq 0$,

$$(4.10) \quad \overline{q}_{x,h}(a) \geq q_{x,h}(a), \text{ for all } a \in \mathbb{R} \text{ and } x \in \overline{T}.$$

Indeed, for any $n \geq 1$,

$$(4.11) \quad \overline{q}_{x,h} \geq 1_{(-\infty,h)} = q_{x,h}^n, \text{ when } x \in \overline{T}_n,$$

and for $x \in \overline{B}_{n-1}$,

$$(4.12) \quad \begin{aligned} q_{x,h}^n &\stackrel{(1.21)}{=} 1_{(-\infty,h)} + 1_{[h,\infty)} \prod_{y^-=x} Q^{\alpha_y}(q_{y,h}^n) \\ &\leq 1_{(-\infty,h)} + 1_{[h,\infty)} \prod_{y^-=x,y \in \overline{T}} Q^{\alpha_y}(q_{y,h}^n) \\ &\leq 1_{(-\infty,h)} + 1_{[h,\infty)} \prod_{y^-=x,y \in \overline{T}} Q^{\overline{\alpha}}(q_{y,h}^n), \end{aligned}$$

where in the last step we have used the fact that $\overline{\alpha} \leq \alpha_y$, for $y \in \overline{T}$, see (4.8), and applied (4.3) of Lemma 4.1 to deduce that $\pi_0 Q^{\overline{\alpha}}(1 - q_{y,h}^n) \leq \pi_0 Q^{\alpha_y}(1 - q_{y,h}^n)$, whence $\pi_h Q^{\alpha_y}(q_{y,h}^n) \leq \pi_h Q^{\overline{\alpha}}(q_{y,h}^n)$ (recall that $h \geq 0$).

Combining (4.11), with (4.9), (4.12), we see that the inequality $\overline{q}_{x,h} \geq q_{x,h}^n$ for $x \in \overline{T}_n$ gets propagated to all $x \in \overline{B}_n$. By Lemma 1.3, letting n tend to infinity we obtain (4.10).

By (1.26) we know that

$$(4.13) \quad \mathbb{P}^G[x_0 \xrightarrow{\varphi \geq h} \infty] = 1 - \mathbb{E}^G[q_{x_0,h}(\varphi_{x_0})].$$

We will show that for small $h > 0$, the right-continuous non-increasing function $\bar{q}_{x_0,h}$ is not identically equal to 1, so that the same holds for $q_{x_0,h}$ by (4.10), and hence the probability in (4.13) is positive. In essence, the proof that $\bar{q}_{x_0,h}$ is not identically equal to 1 will rely on the fact that the largest eigenvalue $\bar{\lambda}_h$ of the Hilbert-Schmidt operator \bar{L}_h defined in (4.16) below, is bigger than 1 for small $h > 0$, see (4.20). This operator is defined on $L^2(\bar{\nu})$, where $\bar{\nu}$ is a centered Gaussian law on \mathbb{R} , which we now introduce.

We denote by $\bar{\nu}$ the centered Gaussian law on \mathbb{R} with variance

$$(4.14) \quad \bar{\sigma}^2 = \frac{\bar{\alpha}}{1 - \bar{\alpha}^2}.$$

Note that (see also (3.10) - (3.16) of [17])

$$(4.15) \quad Q^{\bar{\alpha}} \text{ is a self-adjoint, non-negative Hilbert-Schmidt operator on } L^2(\bar{\nu}).$$

We then consider the self-adjoint, non-negative, Hilbert-Schmidt operator on $L^2(\bar{\nu})$ defined by

$$(4.16) \quad \bar{L}_h = 2\pi_h Q^{\bar{\alpha}} \pi_h, \text{ for } h \in \mathbb{R},$$

as well as its largest eigenvalue (which coincides with its operator norm)

$$(4.17) \quad \bar{\lambda}_h = \|\bar{L}_h\|_{L^2(\bar{\nu}) \rightarrow L^2(\bar{\nu})}.$$

The proof of Proposition 3.1 of [17] shows that

$$(4.18) \quad h \rightarrow \bar{\lambda}_h \text{ is a decreasing homeomorphism from } \mathbb{R} \text{ onto } (0, 2).$$

Moreover, $\pi_0 Q^{\bar{\alpha}} \pi_0(1_{[0,\infty)})(a) = P^Y[\bar{\alpha}a + \sqrt{\bar{\alpha}}Y \geq 0] > \frac{1}{2}$ for $a > 0$, so that

$$(4.19) \quad \bar{\lambda}_0 \geq (\bar{L}_0 1_{[0,\infty)}, 1_{[0,\infty)})_{L^2(\bar{\nu})} / \|1_{[0,\infty)}\|_{L^2(\bar{\nu})}^2 > 1.$$

It thus follows that

$$(4.20) \quad \bar{\lambda}_h > 1 \text{ for small } h > 0.$$

One can then proceed as in the proof of (3.33) of [17] to deduce that

$$(4.21) \quad \bar{\gamma}_h = 1 - \int_h^\infty d\bar{\nu}(a) \bar{q}_{x_0,h}(a) > 0, \text{ when } \bar{\lambda}_h > 1.$$

This shows that for small $h > 0$, $\bar{q}_{x_0,h}$ and hence $q_{x_0,h}$ by (4.10), are not identically equal to 1. As explained below (4.13), it follows that $\mathbb{P}^G[x_0 \xrightarrow{\varphi \geq h} \infty] > 0$ for small $h > 0$, and hence $h_* > 0$. This proves (4.6). \square

Remark 4.4. When M in (4.5) is chosen as an integer bigger or equal to 2, the quantity $\bar{\gamma}_h$ in (4.21) can be interpreted as the probability that the Gaussian free field $\bar{\varphi}$ on an $(M+2)$ -regular tree, when restricted to a given ‘‘forward binary tree’’ rooted at some point, is such that the root belongs to an infinite connected component of $\{\bar{\varphi} \geq h\}$ (see also Section 3 of [17]). \square

As an application of Theorem 3.4 and Proposition 4.2 we have

Corollary 4.5. *Assume that the rooted tree T has bounded degree, contains an infinite binary sub-tree, and $\sup_{x \in T} R_x^\infty < \infty$, then*

$$(4.22) \quad 0 < h_* < \sqrt{2u_*} < \infty.$$

Proof. By the observation below (1.4), moving the root of T if necessary, we can assume that the infinite binary tree is rooted at x_0 , the root of T . The R_x^∞ , $x \in T$, are uniformly bounded, and bounded away from 0 by (1.10) ii). It is now straightforward from (1.34), (1.36) to infer that $0 < u_* < \infty$, see also Theorem 5.1 of [20]. Conditions (3.1) and (3.2) are immediate and the assumptions of Proposition 4.2 are fulfilled as well. The claim (4.22) now follows from Theorem 3.4 and Proposition 4.2. \square

Remark 4.6. In particular, when T is an infinite tree of bounded degree such that outside a finite set, all vertices have degree at least 3, the assumptions of the above corollary are fulfilled by T^∞ , and in view of (1.31), (1.39) one has

$$(4.23) \quad 0 < h_* < \sqrt{2u_*} < \infty.$$

\square

5 Application to super-critical Galton-Watson trees

We will now apply the results of the previous sections to the case where T is a super-critical Galton-Watson tree conditioned on non-extinction, and the root x_0 is the initial ancestor. Our main results appear in Theorems 5.4 and 5.5. An important step is carried out in Proposition 5.2 where it is checked that condition (3.1) holds almost surely. Theorem 5.4 establishes the inequality $h_* < \sqrt{2u_*}$ and the exponential decay in $|x|$ of $\mathbb{P}^G[x_0 \xrightarrow{\varphi \geq 0} x]$ in a broad enough generality. Theorem 5.5 provides a sufficient condition for $h_* > 0$.

We first introduce some notation and recall various known facts. We denote by ν the offspring distribution and assume that

$$(5.1) \quad m = \sum_{k=0}^{\infty} k \nu(k) \in (1, \infty)$$

(we are in the super-critical regime). We denote by P^ν the probability governing the Galton-Watson tree and by f the generating function of ν , which is strictly convex and equals

$$(5.2) \quad f(s) = \sum_{k=0}^{\infty} s^k \nu(k), \quad 0 \leq s \leq 1.$$

If q stands for the extinction probability $P^\nu[|T| < \infty]$, then $q \in [0, 1)$ is the smallest fixed point of f on $[0, 1]$, the only other fixed point being $s = 1$. Moreover, conditioned on non-extinction, i.e. under $P_*^\nu[\cdot] = P^\nu[\cdot \mid |T| = \infty]$, the sub-tree T^∞ of sites with an infinite

line of descent corresponds to a Galton-Watson tree with offspring distribution ν_∞ having the generating function

$$(5.3) \quad f_\infty(s) = \frac{f(q + s(1-q)) - q}{1-q}, \quad 0 \leq s \leq 1,$$

so that $\nu_\infty(0) = 0$, and ν_∞ has same mean m as ν , see for instance Proposition 5.28 of [12]. In addition, one knows that P_*^ν -a.s., T (and T^∞) are transient, see Theorem 3.5 and Corollary 5.10 of [12].

It is further known from Section 3 of [19], that u_* is P_*^ν -a.s. constant and by Theorem 1 and (1.5)' of [19], that

$$(5.4) \quad u_* \in (0, \infty) \text{ is characterized as the unique solution of } f'_\infty(\mathcal{L}(u)) = 1, \text{ where} \\ \mathcal{L}(u) = E_*^\nu[e^{-u(1-\alpha_{x_0})}] (= E^{\nu_\infty}[e^{-u(1-\alpha_{x_0})}]), \text{ for } u \geq 0$$

(E_*^ν, E^{ν_∞} stand for the respective P_*^ν and P^{ν_∞} -expectations).

As we now explain, conditioned on non-extinction, h_* is almost surely constant. We will later see, cf. (5.8), that this constant is finite.

Lemma 5.1. $h_*(T) \geq 0$ is almost everywhere constant under $P_*^\nu + P^{\nu_\infty}$.

Proof. By (1.31) and the observation below (5.2), $h_*(T)$ has same law under P_*^ν and P^{ν_∞} . We thus only need to consider P_*^ν . For $h \in \mathbb{R}$, we introduce the event (see Remark 1.4 for the notation)

$$A_h = \{T; q_{x_0, h}^T \text{ is identically } 1\}.$$

This event (in the space of Galton-Watson trees) is hereditary (or in the terminology of [12], p. 136, the property it describes is inherited). Namely, all finite Galton-Watson trees belong to A_h (by (1.24)), and when $T \in A_h$, the equation

$$q_{x_0, h}^T \stackrel{(1.23)}{=} 1_{(-\infty, h)} + 1_{[h, \infty)} \prod_{|x|=1} Q^{\alpha_x}(q_{x, h}^T)$$

implies that $q_{x, h}^T$ are identically equal to 1 for all $|x| = 1$ in T , and by (1.29) we see that $T_x \in A_h$ for all $|x| = 1$ in T . It now follows by Proposition 5.6 of [12] that $P_*^\nu[A_h] \in \{0, 1\}$ for all $h \in \mathbb{R}$.

By (1.27) we know that for any h in \mathbb{R}

$$(5.5) \quad P_*^\nu\text{-a.s.}, \{h_*(T) < h\} \subseteq A_h \text{ and } \{h_*(T) > h\} \subseteq A_h^c.$$

The random variable $h_*(T)$ is non-negative, see (1.19), so that $P_*^\nu[A_h^c] = 1$ for all $h < 0$. If one sets

$$(5.6) \quad h_0 = \inf\{h \in \mathbb{R}; P_*^\nu[A_h] = 1\} \text{ (with the convention } \inf \phi = \infty),$$

it is now routine to see with (5.5) and the above 0-1 law that P_*^ν -a.s., $h_*(T) = h_0$. This proves Lemma 5.1. \square

We will now see that condition (3.1) is P_*^ν -a.s. fulfilled. In particular, this implies that P_*^ν -a.s. the conclusions of Corollary 2.3 (see Remark 3.1 1)) and of Proposition 3.2 hold.

Proposition 5.2. *There exist $A, M, \delta > 0$ such that P_*^ν -a.s., for large n and all $x \in T^\infty$ with $|x| = n$,*

$$(5.7) \quad \sum_{y \in (x_0, x)} 1\{R_y^\infty \leq A, d_{y^-} \leq M\} \geq \delta n.$$

In particular, the P_^ν -a.s. constant quantities h_* and u_* satisfy*

$$(5.8) \quad 0 \leq h_* \leq \sqrt{2u_*} < \infty.$$

Proof. Once we prove (5.7), the second claim is by Remark 3.1 1) a direct consequence of Corollary 2.3, together with Lemma 5.1 and (5.4). We thus only need to prove (5.7), which pertains to T^∞ . By the observation below (5.2) we can work with P^{ν_∞} in place of P_*^ν , so that $T = T^\infty$, P^{ν_∞} -a.s.. We are going to show that

$$(5.9) \quad \text{for all } \eta > 0, \text{ there exists } M > 0 \text{ such that } P^{\nu_\infty}\text{-a.s., for large } n \text{ and all } x \in T \text{ with } |x| = n, \sum_{y \in [x_0, x]} 1\{d_y \geq M\} \leq \eta n,$$

$$(5.10) \quad \text{there exists } \eta' \in (0, 1) \text{ and } A > 0 \text{ such that } P^{\nu_\infty}\text{-a.s., for large } n \text{ and all } x \in T \text{ with } |x| = n, \sum_{y \in (x_0, x)} 1\{R_y^\infty \leq A\} \geq (1 - \eta')n.$$

Then, choosing $\delta, \eta > 0$ such that $\eta' + \eta + \delta < 1$, it will follow that (5.7) holds with P^{ν_∞} in place of P_*^ν and, as mentioned above, this will prove our claim.

We start with the proof of (5.9). We denote by \bar{P} the measure on Galton-Watson trees endowed with a spine denoted by w_i , $i \geq 0$ (so $w_0 = x_0$ and $w_{i+1}^- = w_i$, for each $i \geq 0$), such that under \bar{P} the individuals on the spine reproduce with the size-biased distribution (recall that ν and ν_∞ have same mean m)

$$(5.11) \quad \bar{\nu}(k) = \frac{k}{m} \nu_\infty(k), \quad k \geq 1,$$

the individuals off the spine reproduce with distribution ν_∞ , and at each step, the next element w_{i+1} on the spine is chosen uniformly among the offspring of w_i (we refer to Chapter 1 §3 of [1], or to Chapter 12 §1 of [12] for details).

Then, for $M, \eta, \lambda > 0$ and $n \geq 1$, we have

$$(5.12) \quad \begin{aligned} & P^{\nu_\infty} \left[\text{for some } x \text{ in } T \text{ with } |x| = n, \sum_{y \in [x_0, x]} 1\{d_y \geq M\} > \eta n \right] \leq \\ & E^{\nu_\infty} \left[\sum_{|x|=n} 1 \left\{ \sum_{y \in [x_0, x]} 1\{d_y \geq M\} > \eta n \right\} \right] = \\ & m^n \bar{P} \left[\sum_{i=0}^{n-1} 1\{d_{w_i} \geq M\} > \eta n \right] \stackrel{\text{exponential Chebyshev}}{\leq} m^n e^{-\lambda \eta n} \bar{E} \left[e^{\lambda \sum_{i=0}^{n-1} 1\{d_{w_i} \geq M\}} \right] \end{aligned}$$

and we made use of Lemma 1.3.2 of [1] for the equality on the third line. The d_{w_i} , $i \geq 0$, are i.i.d. and distributed as $\bar{\nu}$ under \bar{P} . Hence, we have (\bar{E} stands for the \bar{P} -expectation):

$$(5.13) \quad \bar{E}\left[e^{\lambda \sum_{i=0}^{n-1} 1\{d_{w_i} \geq M\}}\right] = \bar{E}\left[e^{\lambda 1\{d_{x_0} \geq M\}}\right]^n \stackrel{(5.11)}{=} m^{-n} (m + (e^\lambda - 1) E^{\nu_\infty}[d_{x_0}, d_{x_0} \geq M])^n.$$

If we now choose $\lambda_0 > 0$ such that

$$(5.14) \quad e^{\frac{\lambda_0}{2} \eta} > m,$$

and M_0 (large) such that

$$(5.15) \quad e^{\frac{\lambda_0}{2} \eta} > m + (e^{\lambda_0} - 1) E^{\nu_\infty}[d_{x_0}, d_{x_0} \geq M_0],$$

we find after insertion in the last line of (5.12) that

$$(5.16) \quad P^{\nu_\infty} \left[\text{for some } x \text{ in } T \text{ with } |x| = n, \sum_{y \in [x_0, x]} 1\{d_y \geq M_0\} > \eta n \right] \leq e^{-\frac{\lambda_0}{2} \eta n}.$$

The claim (5.9) follows by Borel-Cantelli's lemma.

We then turn to the proof of (5.10). We note that if $x \in T$ and $y \in (x_0, x)$, we can set

$$(5.17) \quad \tilde{R}_y = \begin{cases} \infty, & \text{if } d_y = 1, \\ \min\{1 + R_{y'}^\infty; y' \in T_y \text{ and } y' \notin [y, x]\} & \text{otherwise.} \end{cases}$$

Then, clearly $R_y^\infty \leq \tilde{R}_y$. If we now choose A large enough such that $P^{\nu_\infty}[1 + R_{x_0}^\infty \leq A] > 0$, it follows from (2.16) in Lemma 1 of [8], that for some $\eta' \in (0, 1)$

$$(5.18) \quad P^{\nu_\infty} \left[\min_{|x|=n} \sum_{y \in (x_0, x)} 1\{\tilde{R}_y \leq A\} \leq (1 - \eta') n \right] \text{ decays geometrically in } n.$$

By Borel Cantelli's lemma, we see that

$$(5.19) \quad \begin{aligned} & P^{\nu_\infty}\text{-a.s. for large } n \text{ and all } x \in T \text{ with } |x| = n, \\ & \sum_{y \in (x_0, x)} 1\{R_y^\infty \leq A\} \geq \sum_{y \in (x_0, x)} 1\{\tilde{R}_y \leq A\} \geq (1 - \eta') n. \end{aligned}$$

The claim (5.10) follows. This completes the proof of Proposition 5.2. \square

We will now see that the existence of some finite exponential moment of the offspring distribution ν ensures that (3.2) holds P^*_ν -almost surely. This is the last step before Theorem 5.4, which is in essence an application of Theorem 3.4 to the present set-up.

We introduce the condition

$$(5.20) \quad \text{for some } \gamma > 0, \sum_{k \geq 0} e^{\gamma k} \nu(k) < \infty.$$

Under (5.20) the generating function f in (5.2) has an analytic extension to a disc in the complex plane centered at the origin with radius bigger than 1. In view of (5.3) a similar property holds for f_∞ , and hence

$$(5.21) \quad \text{for some } \gamma_\infty > 0, \sum_{k \geq 1} e^{\gamma_\infty k} \nu_\infty(k) < \infty.$$

Proposition 5.3. *When (5.20) holds, then there exists $B > 0$, such that P_*^ν -a.s., for large n and all $x \in T^\infty$ with $|x| = n$,*

$$(5.22) \quad \sum_{y \in (x_0, x]} \frac{1}{R_y^\infty (1 + R_y^\infty)} \leq B n.$$

Proof. As in the proof of Proposition 5.2, we can work with P^{ν_∞} in place of P_*^ν , so that P^{ν_∞} -a.s., $T = T^\infty$. We first observe that for $B > 0$ and $n \geq 1$,

$$(5.23) \quad \begin{aligned} P^{\nu_\infty} \left[\text{for some } x \text{ with } |x| = n, d_x \geq \frac{B}{2} n \right] &\leq m^n P^{\nu_\infty} \left[d_{x_0} \geq \frac{B}{2} n \right] \\ &\leq m^n e^{-\gamma_\infty \frac{B}{2} n} E^{\nu_\infty} [e^{\gamma_\infty d_{x_0}}]. \end{aligned}$$

Moreover, with \bar{P} as below (5.10), and $w_i, i \geq 0$, the spine, we have

$$(5.24) \quad \begin{aligned} P^{\nu_\infty} \left[\text{for some } x \text{ with } |x| = n, \sum_{y \in [x_0, x)} d_y \geq \frac{B}{2} n \right] &\leq \\ E^{\nu_\infty} \left[\sum_{|x|=n} 1 \left\{ \sum_{y \in [x_0, x)} d_y \geq \frac{B}{2} n \right\} \right] &= m^n \bar{P} \left[\sum_{i=0}^{n-1} d_{w_i} \geq \frac{B}{2} n \right]. \end{aligned}$$

Under \bar{P} the variables $d_{w_i}, i \geq 0$, are i.i.d. $\bar{\nu}$ -distributed, with $\bar{\nu}$ the size-biased distribution in (5.11). By the exponential Chernov bound, we find that

$$(5.25) \quad \begin{aligned} \bar{P} \left[\sum_{i=0}^{n-1} d_{w_i} \geq \frac{B}{2} n \right] &\leq \exp \left\{ -n \bar{I} \left(\frac{B}{2} \right) \right\}, \text{ where for } a \geq 0, \\ \bar{I}(a) = \sup_{\lambda \geq 0} \{ \lambda a - \log \bar{E} [e^{\lambda d_{x_0}}] \} &\geq \frac{\gamma_\infty}{2} a - b, \\ \text{with } b = \log \bar{E} [e^{\frac{\gamma_\infty}{2} d_{x_0}}] &(< \infty \text{ by (5.21)}). \end{aligned}$$

If we now choose B_0 large enough so that $\frac{\gamma_\infty}{2} B_0 > \log m$ and $\bar{I}(\frac{B_0}{2}) > \log m$, then we see that the probabilities in (5.23) and (5.24) are summable in n . Hence, by Borel-Cantelli's lemma, we find that P^{ν_∞} -a.s., for large n and all $x \in T$ with $|x| = n$,

$$(5.26) \quad \sum_{y \in (x_0, x]} d_y \leq B_0 n.$$

Since $(R_y^\infty (1 + R_y^\infty))^{-1} \leq d_y$ by (1.10) ii), this proves Proposition 5.3. \square

We now come to one of the main results of this section, which states that for a super-critical Galton-Watson tree conditioned on non-extinction if the offspring distribution has some finite exponential moment then $h_* < \sqrt{2u_*}$. More precisely, one has

Theorem 5.4. *The deterministic critical values h_* and u_* attached to a super-critical Galton-Watson tree conditioned on non-extinction, for which the offspring distribution satisfies (5.1), (5.20), are such that*

$$(5.27) \quad 0 \leq h_* < \sqrt{2u_*} < \infty.$$

In addition, under (5.1) alone, there exists $\beta > 0$ such that P_^ν -a.s.,*

$$(5.28) \quad \text{for large } n \text{ and all } x \in T \text{ with } |x| = n, \mathbb{P}^G [x_0 \overset{\varphi \geq 0}{\longleftrightarrow} x] \leq e^{-\beta n}.$$

Proof. As recalled in (5.4), we know that $0 < u_* < \infty$. Moreover, by Propositions 5.2 and 5.3, we see that P_*^ν -a.s., the conditions (3.1) and (3.2) are fulfilled. The claim (5.27) now follows from Theorem 3.4.

Let us now prove (5.28). By Propositions 5.2 and 3.2 we know that (5.28) holds with T replaced by T^∞ and $e^{-\beta n}$ by $2e^{-\kappa n}$. When the extinction probability $q \in [0, 1)$ vanishes, (5.28) readily follows. Otherwise, we set $m' = f'(q) < 1$ and choose $\eta \in (0, 1)$ so that $\bar{m} = m^{(1-\eta)}m^\eta < 1$. For $n \geq 1$, we set $\tilde{n} = \lceil \eta n \rceil$ and for $x \in T$ such that $|x| = n$, we write \tilde{x} for the site in $[x_0, x]$ such that $|\tilde{x}| = \tilde{n}$. Clearly, $\mathbb{P}^G[x_0 \xrightarrow{\varphi \geq 0} x] \leq \mathbb{P}^G[x_0 \xrightarrow{\varphi \geq 0} \tilde{x}]$, and by the above mentioned exponential decay along T^∞ , the claim (5.28) will follow (with say $\beta = \frac{1}{2}\eta\kappa$) once we show that

$$(5.29) \quad P_*^\nu\text{-a.s. for large } n \text{ and all } x \in T \text{ with } |x| = n, \tilde{x} \in T^\infty.$$

To see this last point, we note that the Galton-Watson tree conditioned on extinction has mean offspring m' , see Lemma 1.2.5 of [1] or Proposition 5.28 of [12], and

$$\begin{aligned} & P^\nu[\text{for some } x \in T \text{ with } |x| = n, \tilde{x} \notin T^\infty] \leq \\ & E^\nu \left[\sum_{|y|=\tilde{n}} (1\{|T_y| < \infty\}) \sum_{|x|=n} 1\{x \in T_y\} \right] = m^{\tilde{n}} q m'^{n-\tilde{n}} \leq q \bar{m}^n. \end{aligned}$$

This last geometric series is convergent and (5.29) follows by Borel Cantelli's lemma. This concludes the proof of Theorem 5.4. \square

We will now introduce a sufficient condition which implies that $h_* > 0$ for the supercritical Galton-Watson tree conditioned on non-extinction under consideration. The argument we use is somewhat in the spirit of [19] for random interacements on Galton-Watson trees, but the situation is more complicated in the case of the Gaussian free field.

Theorem 5.5. *Assume that the mean m of the offspring distribution ν satisfies*

$$(5.30) \quad m > 2,$$

then, almost surely on non-extinction, i.e. P_^ν -a.s.,*

$$(5.31) \quad h_* > 0.$$

Proof. By (1.31) it suffices to consider T^∞ in place of T , and by the observation below (5.2) to replace ν by ν_∞ , so that P^{ν_∞} -a.s., $T^\infty = T$, noting that ν_∞ has the same mean m as ν .

In essence, the strategy of the proof is to obtain for small $h > 0$, uniformly in n , a positive lower bound for the quantity γ_h^n defined below (5.35), which involves the function $r_{x_0, h}^n(\cdot)$. From this lower bound we will infer that P^{ν_∞} -a.s., the function $q_{x_0, h}^n(\cdot)$ is not identically equal to 1, so that $h_* \geq h(> 0)$. The above mentioned lower bound will stem from an inequality based on the branching property (see (5.4) for the definition of $\mathcal{L}(\cdot)$)

$$m' \gamma_h^n > f_{\infty, M}(\mathcal{L}(\frac{h^2}{2})) - f_{\infty, M}(\mathcal{L}(\frac{h^2}{2}) - \gamma_h^n)$$

where $m' \in (2, m)$, and $h > 0$ is such that the derivative of the truncated generating function $f_{\infty, M}$ (see (5.33)) is bigger than m' in a neighborhood of $\mathcal{L}(h^2/2)$.

Let us now proceed with the proof. Our first objective is to establish the key identity that appears in (5.36) below. For $h \in \mathbb{R}$, $n \geq 0$, and x in T with $|x| \leq n$ (from now on we assume that $T^\infty = T$), we consider in the notation of (1.21) the function $r_{x,h}^n(a) = 1 - q_{x,h}^n(a)$, for $a \in \mathbb{R}$, so that

$$\begin{cases} r_{x,h}^n = 1_{[h,\infty)}, \text{ for } x \in T_n, \\ r_{x,h}^n = 1_{[h,\infty)} \left(1 - \prod_{y^-=x} (1 - Q^{\alpha_y}(r_{y,h}^n)) \right), \text{ for } x \text{ in } T, \text{ with } |x| < n \\ \text{(and } r_{x,h}^n \text{ decreases with } n \text{ and tends to } r_{x,h} = 1 - q_{x,h}, \text{ as } n \rightarrow \infty, \\ \text{by Lemma 1.3).} \end{cases}$$

Multiplying both members of the last equality by $e^{-\frac{a^2}{2R_x^\infty}} = \prod_{y^-=x} e^{-\frac{a^2}{2(1+R_y^\infty)}}$, we obtain for x in T with $|x| < n$ and a in \mathbb{R} ,

$$(5.32) \quad e^{-\frac{a^2}{2R_x^\infty}} r_{x,h}^n(a) = 1_{[h,\infty)}(a) \left(\prod_{y^-=x} e^{-\frac{a^2}{2(1+R_y^\infty)}} - \prod_{y^-=x} \left(e^{-\frac{a^2}{2(1+R_y^\infty)}} - e^{-\frac{a^2}{2(1+R_y^\infty)}} Q^{\alpha_y}(r_{y,h}^n(a)) \right) \right).$$

Recall that $(1 + R_{x_0}^\infty)^{-1} = 1 - \alpha_{x_0}$ and $\mathcal{L}(u) = E^{\nu_\infty}[e^{-u(1-\alpha_{x_0})}]$ in the notation of (5.4). For $M \geq 1$, we write

$$(5.33) \quad f_{\infty,M}(s) = \sum_{k \leq M} s^k \nu_\infty(k), \quad 0 \leq s \leq 1.$$

Choosing $x = x_0$, and $n + 1$ in place of n in (5.32), and multiplying both members by the indicator function of the event $\{d_{x_0} \leq M\}$, after P^{ν_∞} -integration and using the branching property, we see that for $h, a \in \mathbb{R}$, $n \geq 0$, and $M \geq 1$

$$(5.34) \quad \begin{aligned} E^{\nu_\infty} \left[e^{-\frac{a^2}{2R_{x_0}^\infty}} r_{x_0,h}^{n+1}(a), d_{x_0} \leq M \right] = \\ 1_{[h,\infty)}(a) \left\{ f_{\infty,M} \left(\mathcal{L} \left(\frac{a^2}{2} \right) \right) - f_{\infty,M} \left(\mathcal{L} \left(\frac{a^2}{2} \right) - E^{\nu_\infty} \left[e^{-\frac{a^2}{2(1+R_{x_0}^\infty)}} Q^{\alpha_{x_0}}(r_{x_0,h}^n(a)) \right] \right) \right\}. \end{aligned}$$

By (5.30) we can now choose a large M , $s_0 \in [0, 1)$, and $m' \in (2, m)$ such that

$$(5.35) \quad f'_{\infty,M}(s) > m', \quad \text{for } s_0 \leq s \leq 1.$$

With this choice of M , when $a = h \geq 0$, we denote by $\tilde{\gamma}_h^n$ the expectation on the left-hand side of (5.34) and by γ_h^n the expectation on the right-hand side so that

$$(5.36) \quad \tilde{\gamma}_h^n = f_{\infty,M} \left(\mathcal{L} \left(\frac{h^2}{2} \right) \right) - f_{\infty,M} \left(\mathcal{L} \left(\frac{h^2}{2} \right) - \gamma_h^n \right), \quad \text{for } n \geq 0 \text{ and } h \geq 0.$$

In addition (since $r_{x_0,h}^n(\cdot)$ is non-decreasing)

$$(5.37) \quad \begin{aligned} \gamma_h^n &\geq e^{-\frac{h^2}{2}} E^{\nu_\infty} [Q^{\alpha_{x_0}}(r_{x_0,h}^n(h))] \stackrel{(1.6)}{\geq} e^{-\frac{h^2}{2}} E^{\nu_\infty} [r_{x_0,h}^n(h) P^Y[\alpha_{x_0} h + \sqrt{\alpha_{x_0}} Y \geq h]] \\ &\geq e^{-\frac{h^2}{2}} E^{\nu_\infty} [d_{x_0} \leq M, r_{x_0,h}^{n+1}(h) \bar{\Phi} \left(\left(\frac{1}{\sqrt{\alpha_{x_0}}} - \sqrt{\alpha_{x_0}} \right) h \right)] > 0, \end{aligned}$$

with $\bar{\Phi}(t) = P^Y[Y > t]$, for $t \in \mathbb{R}$, and we have used that $r_{x_0,h}^{n+1} > 0$ on $[h, \infty)$.

When $\{d_{x_0} \leq M\}$, we have $\alpha_{x_0}^{-1/2} \leq c(M)$ by (1.10), (1.5). Note also that $\bar{\Phi}(0) = 1/2 > 1/m'$. We can thus find by the definition of $\tilde{\gamma}_h^n$ and (5.37) a small $h > 0$, such that

$$(5.38) \quad \gamma_h^n > \frac{1}{m'} \tilde{\gamma}_h^n, \text{ for all } n \geq 0, \text{ as well as } \mathcal{L}\left(\frac{h^2}{2}\right) \geq \frac{1+s_0}{2}.$$

Since $f'_{\infty, M}(s) > m'$ for $s_0 \leq s \leq 1$ it follows from (5.36), (5.38) that $\gamma_h^n \geq \frac{1-s_0}{2}$, for all $n \geq 0$. Hence, by monotone convergence, we find that

$$(5.39) \quad E^{\nu_\infty} \left[e^{-\frac{h^2}{2(1+R_{x_0}^\infty)}} Q^{\alpha_{x_0}}(r_{x_0, h})(h) \right] \geq \frac{1-s_0}{2} > 0.$$

This proves that with positive P^{ν_∞} -measure, $q_{x_0, h}$ is not identically 1, and by the 0-1 law stated above (5.5) this happens P^{ν_∞} -almost surely. By the comment below (5.6) we see that that $h_* \geq h$. This proves (5.31), and concludes the proof of Theorem 5.5. \square

Remark 5.6. Incidentally, Proposition 4.2 implies that $h_* > 0$ for a binary branching, a case corresponding to $m = 2$, which is not covered by Theorem 5.5. One can thus wonder about the nature of broader assumptions under which Theorem 5.5 continues to hold. For instance, does $h_* > 0$ hold almost surely on non-extinction as soon as $m > 1$? \square

A Appendix

In this appendix we provide for the reader's convenience a proof along the argument of Theorem 2 of [2] showing that $h_* \geq 0$ in the general set-up of transient weighted graphs.

We consider a locally finite, connected, transient weighted graph, with vertex set E , and symmetric weights $c_{x,y} = c_{y,x} \geq 0$, which are positive exactly when $x \sim y$, i.e. when x and y are neighbors. We denote by $g(x, y)$, $x, y \in E$, the Green function, by $(\varphi_x)_{x \in E}$ the canonical Gaussian free field, and by \mathbb{P}^G its distribution. The discrete time walk on the weighted graph when located in x jumps to a neighbor y with probability $c_{x,y}/\lambda_x$, where $\lambda_x = \sum_{x' \sim x} c_{x,x'}$. It is governed by the law P_x . We use otherwise similar notation as in Section 1.

We consider a base point $x_0 \in E$. We say that $C \subseteq E$ is a contour surrounding x_0 (in the terminology of [2]), when there exists a finite connected set $K \subseteq E$ containing x_0 such that $C = \partial K$, or when $C = \{x_0\}$ and we set $\{x_0\} = \partial\phi$ by convention. Given a contour C surrounding x_0 , we write $\text{Int } C = K$ for the unique finite connected set K containing x_0 , such that $C = \partial K$, when $C \neq \{x_0\}$, or $K = \phi$, when $C = \{x_0\}$ (when $C \neq \{x_0\}$, $\text{Int } C$ is the connected component of $E \setminus C$ containing x_0).

Given a finite family of contours C_i , $1 \leq i \leq n$ surrounding x_0 , we define the maximal contour via

$$(A.1) \quad \max\{C_1, \dots, C_n\} = \partial\left(\bigcup_{i=1}^n \text{Int } C_i\right),$$

and observe that

$$(A.2) \quad \max\{C_1, \dots, C_n\} \subseteq \bigcup_{i=1}^n C_i.$$

We now consider a finite connected set $U \ni x_0$, $\Delta = \partial U$ and $\bar{U} = U \cup \Delta$. Given $h \in \mathbb{R}$, we introduce the disconnection event

$$(A.3) \quad D_{x_0, U}^h = \{x_0 \text{ is not connected to } \Delta \text{ by a path in } \bar{U} \text{ where } \varphi \geq h\}.$$

Lemma A.1.

$$(A.4) \quad D_{x_0, U}^h = \{\varphi, \text{ there is a contour } C \text{ surrounding } x_0, \text{ with } \\ \text{Int } C \subseteq U, \text{ where } \varphi < h\}.$$

Proof. Denote by \tilde{D} the event on the right-hand side of (A.4). First note that $\tilde{D} \subseteq D_{x_0, U}^h$. Indeed, if C is a contour surrounding x_0 , with $\text{Int } C \subseteq U$ and where $\varphi < h$, any path in \bar{U} from x_0 to $\Delta = \partial U$ will exit $\text{Int } C$ at a point of C where $\varphi < h$. Conversely, one has $D_{x_0, U}^h \subseteq \tilde{D}$. Indeed, when $D_{x_0, U}^h$ occurs, the connected component of $\{\varphi \geq h\}$ containing x_0 is contained in U and its outer boundary (understood as $\{x_0\}$ when this component is empty) yields a contour C with $\text{Int } C \subseteq U$ where $\varphi < h$. This proves the lemma. \square

On the disconnection event $D_{x_0, U}^h$, we can thus define with (A.1)

$$(A.5) \quad C_{<h}^{\max}(U) = \text{the maximal contour of the family of contours } C \\ \text{surrounding } x_0 \text{ with } \text{Int } C \subseteq U, \text{ where } \varphi < h.$$

We recall the definition (0.3) of the critical value h_* .

Proposition A.2.

$$(A.6) \quad h_* \geq 0.$$

Proof. We will show that for any $\varepsilon > 0$,

$$(A.7) \quad \sup_U \mathbb{P}^G[D_{x_0, U}^{-\varepsilon}] < 1$$

(U runs over the collection of finite connected sets containing x_0).

This will imply that for any $\varepsilon > 0$, with positive \mathbb{P}^G -probability the connected component of x_0 in $\{\varphi \geq -\varepsilon\}$ is infinite, and (A.6) will follow.

We thus prove (A.7). For C a contour surrounding x_0 , with $\text{Int } C \subseteq U$, we have by the Markov property of φ under \mathbb{P}^G (see for instance Proposition 2.3 of [18])

$$(A.8) \quad \varphi_{x_0} = h_C + \xi_C \text{ where } h_C = E_{x_0}[\varphi(X_{H_C})] \text{ and } \\ \xi_C \text{ is } N(0, g_{\text{Int } C}(x_0, x_0))\text{-distributed and independent of } \sigma_{(\text{Int } C)^c}$$

(the notation is similar as in (1.7)).

By Lemma A.1 and (A.5), we see that for $h = -\varepsilon$ and $C_{<-\varepsilon}^{\max}$ a shorthand for $C_{<-\varepsilon}^{\max}(U)$

$$(A.9) \quad D_{x_0, U}^{-\varepsilon} = \bigcup_C \{C_{<-\varepsilon}^{\max} = C\}, \text{ where } C \text{ runs over the collection of } \\ \text{contours surrounding } x_0, \text{ with } \text{Int } C \subseteq U.$$

We thus find that

$$(A.10) \quad 0 = \mathbb{E}^G[\text{sign}(\varphi_{x_0})] \stackrel{(A.9)}{=} \sum_C \mathbb{E}^G[\text{sign}(\varphi_{x_0}), C_{<-\varepsilon}^{\max} = C] + \mathbb{E}^G[\text{sign}(\varphi_{x_0}), (D_{x_0,U}^{-\varepsilon})^c],$$

where C runs over the same family as in (A.9).

Note that the event $\{C_{<-\varepsilon}^{\max} = C\}$ is $\sigma_{(\text{Int } C)^c}$ -measurable, and by (A.8) we find that

$$(A.11) \quad \begin{aligned} & \mathbb{E}^G[\text{sign}(\varphi_{x_0}), C_{<-\varepsilon}^{\max} = C] = \mathbb{E}^G[\text{sign}(h_C + \xi_C), C_{<-\varepsilon}^{\max} = C] = \\ & \mathbb{E}^G\left[\left(2\Phi\left(\frac{h_C}{\sqrt{g_{\text{Int } C}(x_0, x_0)}}\right) - 1\right), C_{<-\varepsilon}^{\max} = C\right] \leq \\ & -\left(2\Phi\left(\frac{\varepsilon}{\sqrt{g(x_0, x_0)}}\right) - 1\right) \mathbb{P}^G[C_{<-\varepsilon}^{\max} = C], \end{aligned}$$

where $\Phi(t) = P^Y[Y \leq t]$, for $t \in \mathbb{R}$, with Y a $N(0, 1)$ -distributed variable, and we have used that $h_C \leq -\varepsilon$ on $\{C_{<-\varepsilon}^{\max} = C\}$ and $g_{\text{Int } C}(x_0, x_0) \leq g(x_0, x_0)$ for the last inequality of (A.11).

Coming back to (A.10), we see that

$$(A.12) \quad \begin{aligned} 0 & \leq -\left(2\Phi\left(\frac{\varepsilon}{\sqrt{g(x_0, x_0)}}\right) - 1\right) \sum_C \mathbb{P}^G[C_{<-\varepsilon}^{\max} = C] + 1 - \mathbb{P}^G[D_{x_0,U}^{-\varepsilon}] \\ & \stackrel{(A.9)}{=} -2\Phi\left(\frac{\varepsilon}{\sqrt{g(x_0, x_0)}}\right) \mathbb{P}^G[D_{x_0,U}^{-\varepsilon}] + 1. \end{aligned}$$

This shows that

$$(A.13) \quad \mathbb{P}^G[D_{x_0,U}^{-\varepsilon}] \leq \left(2\Phi\left(\frac{\varepsilon}{\sqrt{g(x_0, x_0)}}\right)\right)^{-1} (< 1),$$

and proves (A.7). Our claim (A.6) follows. \square

Acknowledgements: We thank an anonymous referee for pointing out reference [8], which led to a simplification of the proof of Proposition 5.2.

References

- [1] R. Abraham and J.-F. Delmas. An introduction to Galton-Watson trees and their local limits. Available at <https://hal.archives-ouvertes.fr/hal-01164661>, 2015.
- [2] J. Bricmont, J.L. Lebowitz, and C. Maes. Percolation in strongly correlated systems: the massless Gaussian field. *J. Stat. Phys.*, 48(5/6):1249–1268, 1987.
- [3] J. Černý and A. Teixeira. From random walk trajectories to random interacements. *Ensaaios Matemáticos*, 23, 2012.
- [4] A. Drewitz, B. Ráth, and A. Sapozhnikov. *An Introduction to Random Interacements*. SpringerBriefs in Mathematics, Berlin, 2014.

- [5] A. Drewitz and P.-F. Rodriguez. High-dimensional asymptotics for percolation of Gaussian free field level sets. *Electronic. J. Probab.*, 20(47), 1–39, 2015.
- [6] M. Folz. Volume growth and stochastic completeness of graphs. *Trans. Amer. Math. Soc.*, 366(4):2089–2119, 2014.
- [7] G. Giacomin. Aspects of statistical mechanics of random surfaces. Notes of lectures at IHP (Fall 2001), version of February 24, 2003, available at <http://www.proba.jussieu.fr/pageperso/giacomin/pub/IHP.ps>, 2003.
- [8] G. Grimmett and H. Kesten. Random electrical networks on complete graphs II: Proofs. Available at arXiv:math/0107068.
- [9] T. Lupu. From loop clusters and random interlacement to the free field. To appear in *Ann. Probab.*, available at arXiv:1402.0298.
- [10] R. Lyons. Random walks and percolation on trees. *Ann. Probab.*, 18(3):931–958, 1990.
- [11] R. Lyons. Random walks, capacity and percolation on trees. *Ann. Probab.*, 20(4):2043–2088, 1992.
- [12] R. Lyons and Y. Peres. *Probability on Trees and Networks*. Available at <http://pages.iu.edu/~rdlyons/>, 2016.
- [13] P.-F. Rodriguez and A.S. Sznitman. Phase transition and level-set percolation for the Gaussian free field. *Commun. Math. Phys.*, 320:571–601, 2013.
- [14] J. Rosen. Lectures on Isomorphism Theorems. *Preprint*, available at arXiv:1407.1559.
- [15] C. Sabot and P. Tarrès. Inverting Ray-Knight identity. To appear in *Probab. Theory Relat. Fields*, available at arXiv:1311.6622.
- [16] A.S. Sznitman. An isomorphism theorem for random interacements. *Electron. Commun. Probab.*, 17(9):1–9, 2012.
- [17] A.S. Sznitman. Coupling and an application to level-set percolation of the Gaussian free field. *Electron. J. Probab.*, 21(35), 1–26, 2016 .
- [18] A.S. Sznitman. Topics in occupation times and Gaussian free fields. *Zurich Lectures in Advanced Mathematics*, EMS, Zurich, 2012.
- [19] M. Tassy. Random interacements on Galton-Watson trees. *Electron. Commun. Probab.*, 15:562–571, 2010.
- [20] A. Teixeira. Interlacement percolation on transient weighted graphs. *Electron. J. Probab.*, 14:1604–1627, 2009.
- [21] A. Zhai. Exponential concentration of cover times. *Preprint*, available at arXiv:1407.7617.