

Portfolio Optimisation under Transaction Costs

W. Schachermayer

University of Vienna
Faculty of Mathematics

joint work with
Ch. Czichowsky (Univ. Vienna), J. Muhle-Karbe (ETH Zürich)

September 2012

We fix a strictly positive càdlàg stock price process $S = (S_t)_{0 \leq t \leq T}$.

For $0 < \lambda < 1$ we consider the bid-ask spread $[(1 - \lambda)S, S]$.

A self-financing trading strategy is a predictable, finite variation process $\varphi = (\varphi_t^0, \varphi_t^1)_{0 \leq t \leq T}$ such that

$$d\varphi_t^0 \leq -S_t(d\varphi_t^1)_+ + (1 - \lambda)S_t(d\varphi_t^1)_-$$

φ is called 0-admissible if

$$\varphi_t^0 + (1 - \lambda)S_t(\varphi_t^1)_+ - S_t(\varphi_t^1)_- \geq 0$$

We fix a strictly positive càdlàg stock price process $S = (S_t)_{0 \leq t \leq T}$.

For $0 < \lambda < 1$ we consider the bid-ask spread $[(1 - \lambda)S, S]$.

A self-financing trading strategy is a predictable, finite variation process $\varphi = (\varphi_t^0, \varphi_t^1)_{0 \leq t \leq T}$ such that

$$d\varphi_t^0 \leq -S_t(d\varphi_t^1)_+ + (1 - \lambda)S_t(d\varphi_t^1)_-$$

φ is called 0-admissible if

$$\varphi_t^0 + (1 - \lambda)S_t(\varphi_t^1)_+ - S_t(\varphi_t^1)_- \geq 0$$

Definition [Jouini-Kallal ('95), Cvitanic-Karatzas ('96), Kabanov-Stricker ('02),...]

A *consistent-price system* is a pair (\tilde{S}, Q) such that $Q \sim \mathbb{P}$, the process \tilde{S} takes its value in $[(1 - \lambda)S, S]$, and \tilde{S} is a Q -martingale.

Identifying Q with its density process

$$Z_t^0 = \mathbb{E} \left[\frac{dQ}{d\mathbb{P}} \middle| \mathcal{F}_t \right], \quad 0 \leq t \leq T$$

we may identify (\tilde{S}, Q) with the \mathbb{R}^2 -valued martingale $Z = (Z_t^0, Z_t^1)_{0 \leq t \leq T}$ such that

$$\tilde{S} := \frac{Z_t^1}{Z_t^0} \in [(1 - \lambda)S, S].$$

For $0 < \lambda < 1$, we say that S satisfies (CPS^λ) if there is a consistent price system for transaction costs λ .

Definition [Jouini-Kallal ('95), Cvitanic-Karatzas ('96), Kabanov-Stricker ('02),...]

A *consistent-price system* is a pair (\tilde{S}, Q) such that $Q \sim \mathbb{P}$, the process \tilde{S} takes its value in $[(1 - \lambda)S, S]$, and \tilde{S} is a Q -martingale.

Identifying Q with its density process

$$Z_t^0 = \mathbb{E} \left[\frac{dQ}{d\mathbb{P}} \middle| \mathcal{F}_t \right], \quad 0 \leq t \leq T$$

we may identify (\tilde{S}, Q) with the \mathbb{R}^2 -valued martingale $Z = (Z_t^0, Z_t^1)_{0 \leq t \leq T}$ such that

$$\tilde{S} := \frac{Z_t^1}{Z_t^0} \in [(1 - \lambda)S, S].$$

For $0 < \lambda < 1$, we say that S satisfies (CPS^λ) if there is a consistent price system for transaction costs λ .

Remark [Guasoni, Rasonyi, S. ('08)]

If the process $S = (S_t)_{0 \leq t \leq T}$ is *continuous* and has *conditional full support*, then (CPS^μ) is satisfied, for all $\mu > 0$.

For example, exponential fractional Brownian motion verifies this property.

Portfolio optimisation

The set of non-negative claims attainable at price x is

$$\mathcal{C}(x) = \left\{ \begin{array}{l} X_T \in L_+^0 : \text{there is a 0-admissible } \varphi = (\varphi_t^0, \varphi_t^1)_{0 \leq t \leq T} \\ \text{starting at } (\varphi_0^0, \varphi_0^1) = (x, 0) \text{ and ending at} \\ (\varphi_T^0, \varphi_T^1) = (X_T, 0) \end{array} \right\}$$

Given a utility function $U : \mathbb{R}_+ \rightarrow \mathbb{R}$ define

$$u(x) = \sup\{\mathbb{E}[U(X_T)] : X_T \in \mathcal{C}(x)\}.$$

Cvitanic-Karatzas ('96), Deelstra-Pham-Touzi ('01),
Cvitanic-Wang ('01), Bouchard ('02),...

Question 1

What are conditions ensuring that $\mathcal{C}(x)$ is closed in $L^0_+(\mathbb{P})$. (w.r. to convergence in measure) ?

Theorem [Cvitanic-Karatzas ('96), Campi-S. ('06)]:

Suppose that (CPS^μ) is satisfied, for all $\mu > 0$, and fix $\lambda > 0$.
Then $\mathcal{C}(x) = \mathcal{C}^\lambda(x)$ is closed in L^0 .

Theorem [Guasoni, Rasonyi, S. ('08)]:

Let $S = (S_t)_{0 \leq t \leq T}$ be a continuous process. TFAE

- (i) For each $\mu > 0$, S does not allow for arbitrage under transaction costs μ .
- (ii) For each $\mu > 0$, (CPS^μ) holds, i.e. *consistent price systems under transaction costs μ* exist.

Question 1

What are conditions ensuring that $\mathcal{C}(x)$ is closed in $L^0_+(\mathbb{P})$. (w.r. to convergence in measure) ?

Theorem [Cvitanic-Karatzas ('96), Campi-S. ('06)]:

Suppose that (CPS^μ) is satisfied, for all $\mu > 0$, and fix $\lambda > 0$. Then $\mathcal{C}(x) = \mathcal{C}^\lambda(x)$ is closed in L^0 .

Theorem [Guasoni, Rasonyi, S. ('08)]:

Let $S = (S_t)_{0 \leq t \leq T}$ be a continuous process. TFAE

- (i) For each $\mu > 0$, S does not allow for arbitrage under transaction costs μ .
- (ii) For each $\mu > 0$, (CPS^μ) holds, i.e. *consistent price systems under transaction costs μ* exist.

Question 1

What are conditions ensuring that $\mathcal{C}(x)$ is closed in $L^0_+(\mathbb{P})$. (w.r. to convergence in measure) ?

Theorem [Cvitanic-Karatzas ('96), Campi-S. ('06)]:

Suppose that (CPS^μ) is satisfied, for all $\mu > 0$, and fix $\lambda > 0$. Then $\mathcal{C}(x) = \mathcal{C}^\lambda(x)$ is closed in L^0 .

Theorem [Guasoni, Rasonyi, S. ('08)]:

Let $S = (S_t)_{0 \leq t \leq T}$ be a continuous process. TFAE

- (i) For each $\mu > 0$, S does not allow for arbitrage under transaction costs μ .
- (ii) For each $\mu > 0$, (CPS^μ) holds, i.e. *consistent price systems under transaction costs μ exist.*

The dual objects

Definition

We denote by $D(y)$ the convex subset of $L_+^0(\mathbb{P})$

$$D(y) = \{yZ_T^0 = y \frac{dQ}{d\mathbb{P}}, \text{ for some consistent price system } (\tilde{S}, Q)\}$$

and

$$\mathcal{D}(y) = \overline{\text{sol}(D(y))}$$

the closure of the solid hull of $D(y)$ taken with respect to convergence in measure.

Definition [Kramkov-S. ('99), Karatzas-Kardaras ('06), Campi-Owen ('11),...]

We call a process $Z = (Z_t^0, Z_t^1)_{0 \leq t \leq T}$ a *super-martingale deflator* if $Z_0^0 = 1$, $\frac{Z_t^1}{Z_t^0} \in [(1 - \lambda)S, S]$, and for each 0-admissible, self-financing φ the value process

$$\varphi_t^0 Z_t^0 + \varphi_t^1 Z_t^1 = Z_t^0 (\varphi_t^0 + \varphi_t^1 \frac{Z_t^1}{Z_t^0})$$

is a super-martingale.

Proposition

$$\mathcal{D}(y) = \{yZ_T^0 : Z = (Z_t^0, Z_t^1)_{0 \leq t \leq T} \text{ a super - martingale deflator}\}$$

Definition [Kramkov-S. ('99), Karatzas-Kardaras ('06), Campi-Owen ('11),...]

We call a process $Z = (Z_t^0, Z_t^1)_{0 \leq t \leq T}$ a *super-martingale deflator* if $Z_0^0 = 1$, $\frac{Z_t^1}{Z_t^0} \in [(1 - \lambda)S, S]$, and for each 0-admissible, self-financing φ the value process

$$\varphi_t^0 Z_t^0 + \varphi_t^1 Z_t^1 = Z_t^0 (\varphi_t^0 + \varphi_t^1 \frac{Z_t^1}{Z_t^0})$$

is a super-martingale.

Proposition

$\mathcal{D}(y) = \{yZ_T^0 : Z = (Z_t^0, Z_t^1)_{0 \leq t \leq T} \text{ a super - martingale deflator}\}$

Theorem (Czichowsky, Muhle-Karbe, S. ('12))

Let S be a càdlàg process, $0 < \lambda < 1$, suppose that (CPS^μ) holds true, for each $\mu > 0$, suppose that U has reasonable asymptotic elasticity and $u(x) < U(\infty)$, for $x < \infty$.

Then $\mathcal{C}(x)$ and $\mathcal{D}(y)$ are polar sets:

$$X_T \in \mathcal{C}(x) \quad \text{iff} \quad \langle X_T, Y_T \rangle \leq xy, \quad \text{for } Y_T \in \mathcal{D}(y)$$

$$Y_T \in \mathcal{D}(y) \quad \text{iff} \quad \langle X_T, Y_T \rangle \leq xy, \quad \text{for } X_T \in \mathcal{C}(y)$$

Therefore by the abstract results from [Kramkov-S. ('99)] the duality theory for the portfolio optimisation problem works as nicely as in the frictionless case: for $x > 0$ and $y = u'(x)$ we have

(i) There is a unique primal optimiser $\hat{X}_T(x) = \hat{\varphi}_T^0$
which is the terminal value of an optimal $(\hat{\varphi}_t^0, \hat{\varphi}_t^1)_{0 \leq t \leq T}$.

(i') There is a unique dual optimiser $\hat{Y}_T(y) = \hat{Z}_T^0$
which is the terminal value of an optimal
super-martingale deflator $(\hat{Z}_t^0, \hat{Z}_t^1)_{0 \leq t \leq T}$.

(ii) $U'(\hat{X}_T(x)) = \hat{Z}_t^0(y), \quad -V'(\hat{Z}_T(y)) = \hat{X}_T(x)$

(iii) The process $(\hat{\varphi}_t^0 \hat{Z}_t^0 + \hat{\varphi}_t^1 \hat{Z}_t^1)_{0 \leq t \leq T}$ is a martingale, and
therefore

$$\{d\hat{\varphi}_t^0 > 0\} \subseteq \left\{ \frac{\hat{Z}_t^1}{\hat{Z}_t^0} = (1 - \lambda)S_t \right\},$$

$$\{d\hat{\varphi}_t^0 < 0\} \subseteq \left\{ \frac{\hat{Z}_t^1}{\hat{Z}_t^0} = S_t \right\},$$

etc. etc.

Theorem [Cvitanic-Karatzas ('96)]

In the setting of the above theorem *suppose* that $(\hat{Z}_t)_{0 \leq t \leq T}$ is a local martingale.

Then $\hat{S} = \frac{\hat{Z}_1}{\hat{Z}_0}$ is a *shadow price*, i.e. the optimal portfolio for the *frictionless market* \hat{S} and for the *market* S under *transaction costs* λ coincide.

Sketch of Proof

Suppose (w.l.g.) that $(\hat{Z}_t)_{0 \leq t \leq T}$ is a true martingale. Then $\frac{d\hat{Q}}{d\mathbb{P}} = \hat{Z}_T^0$ defines a *probability measure* under which the process $\hat{S} = \frac{\hat{Z}_1}{\hat{Z}_0}$ is a martingale. Hence we may apply the frictionless theory to (\hat{S}, \mathbb{P}) .

\hat{Z}_T^0 is (a fortiori) the dual optimizer for \hat{S} .

As \hat{X}_T and \hat{Z}_T^0 satisfy the first order condition

$$U'(\hat{X}_T) = \hat{Z}_T^0,$$

\hat{X}_T must be the optimizer for the frictionless market \hat{S} too. ■

Theorem [Cvitanic-Karatzas ('96)]

In the setting of the above theorem *suppose* that $(\hat{Z}_t)_{0 \leq t \leq T}$ is a local martingale.

Then $\hat{S} = \frac{\hat{Z}_1}{\hat{Z}_0}$ is a *shadow price*, i.e. the optimal portfolio for the *frictionless market* \hat{S} and for the *market* S under *transaction costs* λ coincide.

Sketch of Proof

Suppose (w.l.g.) that $(\hat{Z}_t)_{0 \leq t \leq T}$ is a true martingale. Then $\frac{d\hat{Q}}{d\mathbb{P}} = \hat{Z}_T^0$ defines a *probability measure* under which the process $\hat{S} = \frac{\hat{Z}_1}{\hat{Z}_0}$ is a martingale. Hence we may apply the frictionless theory to (\hat{S}, \mathbb{P}) .

\hat{Z}_T^0 is (a fortiori) the dual optimizer for \hat{S} .

As \hat{X}_T and \hat{Z}_T^0 satisfy the first order condition

$$U'(\hat{X}_T) = \hat{Z}_T^0,$$

\hat{X}_T must be the optimizer for the frictionless market \hat{S} too. ■

Question

When is the dual optimizer \hat{Z} a *local martingale*?
Are there cases when it only is a *super-martingale*?

Theorem [Czichowsky-S. ('12)]

Suppose that S is *continuous* and satisfies (*NFLVR*), and suppose that U has reasonable asymptotic elasticity. Fix $0 < \lambda < 1$ and suppose that $u(x) < U(\infty)$, for $x < \infty$.

Then the dual optimizer \hat{Z} is a local martingale. Therefore $\hat{S} = \frac{\hat{Z}^1}{\hat{Z}^0}$ is a shadow price.

Remark

The condition (*NFLVR*) cannot be replaced by requiring (*CPS $^\lambda$*), for each $\lambda > 0$.

Theorem [Czichowsky-S. ('12)]

Suppose that S is *continuous* and satisfies (*NFLVR*), and suppose that U has reasonable asymptotic elasticity. Fix $0 < \lambda < 1$ and suppose that $u(x) < U(\infty)$, for $x < \infty$.

Then the dual optimizer \hat{Z} is a local martingale. Therefore $\hat{S} = \frac{\hat{Z}^1}{\hat{Z}^0}$ is a shadow price.


Remark

The condition (*NFLVR*) cannot be replaced by requiring (*CPS $^\lambda$*), for each $\lambda > 0$.

Examples

Frictionless Example [Kramkov-S. ('99)]

Let $U(x) = \log(x)$. The stock price $S = (S_t)_{t=0,1}$ is given by



Here $\sum_{n=1}^{\infty} \varepsilon_n = 1 - p \ll 1$.


For $x = 1$ the optimal strategy is to buy one stock at time 0 i.e. $\hat{\varphi}_1^1 = 1$.

Let $A_n = \{S_1 = \frac{1}{n}\}$ and consider $A_{\infty} = \{S_1 = 0\}$ so that $\mathbb{P}[A_n] = \varepsilon_n > 0$, for $n \in \mathbb{N}$, while $\mathbb{P}[A_{\infty}] = 0$.

Intuitively speaking, the constraint $\hat{\varphi}_1^1 \leq 1$ comes from the null-set A_{∞} rather than from any of the A_n 's.

It turns out that the dual optimizer \hat{Z} verifies $\mathbb{E}[\hat{Z}_1] < 1$, i.e. only is a super-martingale. Intuitively speaking, the optimal measure \hat{Q} gives positive mass to the \mathbb{P} -null set A_{∞} (compare Cvitanic-Schachermayer-Wang ('01), Campi-Owen ('11)).

Discontinuous Example under transaction costs λ
 (Czichowsky, Muhle-Karbe, S. ('12), compare also Benedetti, Campi, Kallsen, Muhle-Karbe ('11)).



For $x = 1$ it is optimal to buy $\frac{1}{1+\lambda}$ many stocks at time 0. Again, the constraint comes from the \mathbb{P} -null set $A_\infty = \{S_1 = 1\}$.

There is no shadow-price. The intuitive reason is again that the binding constraint on the optimal strategy comes from the \mathbb{P} -null set $A_\infty = \{S_1 = 1\}$.

Continuous Example under Transaction Costs [Czichowsky-S. ('12)]

Let $(W_t)_{t \geq 0}$ be a Brownian motion, starting at $W_0 = w > 0$, and

$$\tau = \inf\{t : W_t - t \leq 0\}$$

Define the stock price process

$$S_t = e^{t \wedge \tau}, \quad t \geq 0.$$

S does not satisfy (*NFLVR*), but it does satisfy (*CPS* $^\lambda$), for all $\lambda > 0$.

Fix $U(x) = \log(x)$, transaction costs $0 < \lambda < 1$, and the initial endowment $(\varphi_0^0, \varphi_0^1) = (1, 0)$.

For the trade at time $t = 0$, we find three regimes determined by thresholds $0 < \underline{w} < \bar{w} < \infty$.

- (i) if $w \leq \underline{w}$ we have $(\hat{\varphi}_{0+}^0, \hat{\varphi}_{0+}^1) = (1, 0)$, i.e. no trade.
- (ii) if $\underline{w} < w < \bar{w}$ we have $(\hat{\varphi}_{0+}^0, \hat{\varphi}_{0+}^1) = (1 - a, a)$, for some $0 < a < \frac{1}{\lambda}$.
- (iii) if $w \geq \bar{w}$, we have $(\hat{\varphi}_{0+}^0, \hat{\varphi}_{0+}^1) = (1 - \frac{1}{\lambda}, \frac{1}{\lambda})$, so that the liquidation value is zero (maximal leverage).

We now choose $W_0 = w$ with $w > \bar{w}$.

Note that the optimal strategy $\hat{\varphi}$ *continues to increase the position in stock*, as long as $W_t - t \geq \bar{w}$.

If there were a shadow price \hat{S} , we therefore necessarily would have

$$\hat{S}_t = e^t, \quad \text{for } 0 \leq t \leq \inf\{u : W_u - u \leq \bar{w}\}.$$

But this is absurd, as \hat{S} clearly does not allow for an e.m.m.

Problem

Let $(B_t^H)_{0 \leq t \leq T}$ be a fractional Brownian motion with Hurst index $H \in]0, 1[\setminus \{\frac{1}{2}\}$. Let $S = \exp(B_t^H)$, and fix $\lambda > 0$ and $U(x) = \log(x)$.

Is the dual optimiser a local martingale or only a super-martingale?
Equivalently, is there a shadow price \hat{S} ?